

# Metal-coupled folding as the driving force for the extreme stability of Rad50 zinc hook dimer assembly

Tomasz Kochańczyk,<sup>1</sup> Michał Nowakowski,<sup>2</sup> Dominika Wojewska,<sup>1</sup> Anna Kocyła,<sup>1</sup> Andrzej Ejchart,<sup>3</sup> Wiktor Koźmiński<sup>2</sup> & Artur Krężel\*<sup>1</sup>

Contribution from

<sup>1</sup> *Department of Chemical Biology, Faculty of Biotechnology, University of Wrocław, Joliot-Curie 14a, 50-383 Wrocław, Poland*

<sup>2</sup> *Biological and Chemical Research Center, Faculty of Chemistry, University of Warsaw, Żwirki i Wigury 101, 02-089 Warsaw, Poland*

<sup>3</sup> *Institute of Biochemistry and Biophysics, Polish Academy of Sciences, Pawińskiego 5a, 02-106, Warsaw, Poland*

\* Corresponding author, E-mail: artur.krezel@uwr.edu.pl

## Abstract

The binding of metal ions at the interface of protein complexes presents a unique and poorly understood mechanism of molecular assembly. A remarkable example is the Rad50 zinc hook domain, which is highly conserved and facilitates the  $\text{Zn}^{2+}$ -mediated homodimerization of Rad50 proteins. Here, we present a detailed analysis of the structural and thermodynamic effects governing the formation and stability ( $\log K_{12} = 20.74$ ) of this evolutionarily conserved protein assembly. We have dissected the determinants of the stability contributed by the small  $\beta$ -hairpin of the domain surrounding the zinc binding motif and the coiled-coiled regions using peptides of various lengths from 4 to 45 amino acid residues, alanine substitutions and peptide bond-to-ester perturbations. In the studied series of peptides, an  $>650\,000$ -fold increase of the formation constant of the dimeric complex arises from favorable enthalpy because of the increased acidity of the cysteine thiols in metal-free form and the structural properties of the dimer. The dependence of the enthalpy on the domain fragment length is partially compensated by the entropic penalty of domain folding, indicating enthalpy-entropy compensation. This study facilitates understanding of the metal-mediated protein-protein interactions in which the metal ion is critical for the tight association of protein subunits.

## Introduction

Transition metal ions play an important role in facilitating the diverse functions of proteins. Among all transition metal ions, zinc (formally  $\text{Zn}^{2+}$ ) is the most widespread cofactor in proteins<sup>1,2</sup>. Based on their function, the zinc binding domains in proteins have been categorized into catalytic, structural and regulatory classes. Another classification scheme is based on the architecture of particular zinc binding domains, namely, the number of polypeptide chains or protein subunits that contribute to  $\text{Zn}^{2+}$  binding. The vast majority of known zinc domains, such as the zinc fingers and catalytic domains in enzymes, have an intramolecular binding architecture, in which all protein-derived zinc ligands are located within a single polypeptide chain<sup>1,2</sup>. Alternatively,  $\text{Zn}^{2+}$  can be intermolecularly bound, i.e., by two or more peptide chains, bridging these molecules to form a protein assembly<sup>3-5</sup>, as in the case of the Rad50 dimer protein<sup>6</sup> or a higher-order assembly<sup>7,8</sup>.

Rad50 protein is a member of an evolutionarily conserved Mre11 complex (Mre11, Rad50, and Nbs1 in eukaryotes) that plays a pivotal role in the DNA damage response, including cell cycle checkpoint activation in response to DNA double-strand breaks (DSBs), DSB repair and telomere maintenance<sup>9</sup>. Rad50 is a rod-shaped molecule with a long antiparallel coiled-coil protruding from the globular ATPase/DNA-binding domain. The apex of the coiled-coil contains a small loop with a conserved Cys-Xaa-Xaa-Cys (CXXC) motif. Rad50 forms a homodimer through two pairs of cysteines from CXXC motifs that contribute to form an intermolecular zinc complex known as zinc hook domain (Fig. 1a). The zinc hook is conserved in Rad50 homologs in all forms of life, from Archaea to humans and even viruses<sup>10</sup>. The global influence of the zinc hook on the functions of Rad50 is universal as disruption of the  $\text{Zn}^{2+}$  binding to the zinc hook is lethal<sup>11,12</sup>. The destabilization of the zinc hook

complex has a long-range allosteric effect on the several hundred angstroms distant globular domain of Rad50, affecting the DNA damage response, DNA recombination, telomere integrity and meiosis. The precise role of the zinc hook complex in these processes and how the stability of this complex relates to its function are unknown<sup>11,13</sup>.

Several intramolecular  $\text{Zn}^{2+}$  binding domains, such as zinc fingers, have been studied in considerable detail regarding metal-binding affinity, thermodynamics and structure-stability relationships<sup>14,15</sup>. In contrast, there is a lack of analyses aimed at understanding the energetic and structural effects accompanying the formation of protein-interface  $\text{Zn}^{2+}$  binding domains, although this aspect is crucial for understanding the roles of such domains in biological processes<sup>16</sup>.

Specific protein-protein interactions are typically stabilized through an extensive network of non-covalent interactions that outcompete the energetic cost attributed to the loss of entropy and the dehydration of protein surfaces<sup>17,18</sup>. The binding of metal ions at the protein-protein interface significantly contributes to the complex formation thermodynamics, and may promote molecular recognition and induce changes in the conformation of the resulting complex. To achieve sufficient stability and specificity, the protein-protein interface containing only non-covalent interactions must bury a large surface area of typically 1500 to 10 000 Å<sup>2</sup><sup>16,19</sup>. The total contact area of the interface formed by the zinc hook domain of Rad50 protein is only 640 Å<sup>2</sup>,<sup>6,20</sup> suggesting a major effect from a low number of interactions and a significant contribution of intermolecular  $\text{Zn}^{2+}$  binding to the stability of the interaction.

Based on the currently available structure of the zinc hook domain of Rad50 from *P. furiosus* (PDB ID: 1L8D)<sup>6</sup> (Fig. 1a), we designed a series of peptides of various lengths ranging from the CXXC motif required for intermolecular  $\text{Zn}^{2+}$  binding

to the full-length domain. We also used alanine substitutions and the replacement of the amide bond with an ester bond to address the influence of certain residues and hydrogen bond formation in the peptide backbone on zinc hook complex stability. Using these models, we dissected the structural and thermodynamic determinants governing the stability of the metal-mediated dimer assembly formed by the zinc hook domain.

## Results and discussion


Based on the previous studies<sup>6</sup> demonstrating the absolute requirement of  $\text{Zn}^{2+}$  for formation of the Rad50 dimer, we envisioned the formation of the complex as a reaction between two monomers and the  $\text{Zn}^{2+}$  ion to form a homodimeric zinc hook complex,  $\text{Zn}(\text{Hk})_2$ . For the structural and thermodynamic analysis of the zinc hook domain, we designed a series of zinc hook (Hk) peptides ranging in length from 4 to 45 amino acids (Hk4-Hk45). The 45-amino acid zinc hook domain peptide model (Hk45) used in the present study includes all residues that form intermolecular contacts in the complex and contains both the  $\alpha$ -helical and  $\beta$ -hairpin regions of the domain (Fig. 1b, Fig. 1c)<sup>20</sup>. To examine the effect of the helical region on the  $\text{Zn}^{2+}$  binding thermodynamics, we shortened the helical region from both the N- and C-termini to 37 (Hk37), 31 (Hk31), 27 (Hk27) and 23 (Hk23) amino acid residues (Fig. 1d). Similarly, we designed a series of peptides (Hk4-Hk14) to investigate the structural and energetic effects of the formation of the central  $\beta$ -hairpin in the zinc hook structure (Fig. 1e).

**$\text{Zn}^{2+}$ -coupled folding of the hook domain.** The binding of  $\text{Zn}^{2+}$  ions to hook peptides was monitored by circular dichroism spectroscopy (CD). Far-UV CD titrations, performed by the addition of  $\text{Zn}^{2+}$  to the initially metal-free peptide at pH 7.4 in the


presence of TCEP (non-metal binding reductant)<sup>21</sup>, demonstrated extensive conformational changes upon  $\text{Zn}^{2+}$  binding. As expected, all hook peptides preferentially form  $\text{Zn}^{2+}$ -mediated dimers ( $\text{Zn}(\text{Hk})_2$ ) at pH 7.4, as indicated by a sharp inflection in the titration curves at a 1:2  $\text{Zn}^{2+}$ -to-peptide molar ratio (insets of Fig. 2a and Fig. S1). Depending on the examined peptide, the  $\text{Zn}(\text{Hk})_2$  complexes adopt different conformations, as indicated by the distinct CD spectra obtained (blue curves, Fig. 2a, Fig. S1). The structural changes occurring during the association of hook subunits into  $\text{Zn}(\text{Hk})_2$  can be compared based on the differential spectra obtained through the spectral subtraction of the hook peptides from their  $\text{Zn}^{2+}$  complexes (Fig. 2b). The differential spectra of the Hk4-Hk14 series showed features characteristic for the CD spectra of  $\beta$ -rich proteins and isolated type II  $\beta$ -turns<sup>22</sup>. The differential CD spectra of the Hk23-Hk27 series were similar to those of the Hk10-Hk14 series, likely reflecting the fact that the helical fragments of the domains in Hk23 and Hk27 are too short to form the stable helical/coiled-coil structure observed in the crystal structure. For the Hk31-Hk45 series, as the peptide length increased, a gradual increase in negative ellipticity and an increase in the ratio of ellipticities at 208 and 222 nm were observed, indicating an increase in the helical and coiled-coil content in the structure<sup>23</sup>. This increase in  $\text{Zn}^{2+}$ -dependent structural changes with peptide length indicates that  $\text{Zn}^{2+}$  binding is coupled to the global folding of the domain, which potentially propagates to the  $\sim 500$  Å distant globular domain of Rad50<sup>13</sup>. This interpretation is consistent with previous findings showing that the disruption of  $\text{Zn}^{2+}$  binding to the hook domain results in the complete loss of Rad50 activity<sup>6,24</sup> and that mutations affecting the structure of the zinc hook influence functions in the globular domain of Rad50<sup>11,13</sup>. Interestingly, quite similar  $\text{Zn}^{2+}$ -coupled folding associated with

cooperative formation of a helical motif was observed for model peptides of a treble-clef zinc finger<sup>25</sup>.

**Stability of the zinc hook domain.** We used potentiometric (Hk4-Hk14) and competitive CD titrations with zinc chelators (Hk23-Hk45) to measure the stability of the zinc hook complexes (Table 1, Table S1, Fig. S3, Fig. S4)<sup>15,26</sup>. Both potentiometric and CD titrations of Hk peptides with  $\text{Zn}^{2+}$  indicated that  $\text{Zn}^{2+}$ -mediated dimers ( $\text{Zn}(\text{Hk})_2$ ) are formed preferentially at pH 7.4 (Fig. 2, Fig. S3). The cumulative constant ( $K_{12}$ , also known as  $\beta_2$ ) for the formation of the dimeric  $\text{Zn}(\text{Hk})_2$  complex is given by equation (1):



The stepwise constants  $K_1$  and  $K_2$  refer to the formation of the complexes one step at a time as noted in equations (2) and (3):



The apparent formation constants ( $\log K_{12}$ ) of the dimeric  $\text{Zn}(\text{Hk})_2$  complexes at pH 7.4, calculated either from protonation and stability constants determined potentiometrically (see example in Supplementary information) or from CD competition experiments with zinc chelators, are presented in Table 1. It should be noted that the formation constant of  $\text{Zn}(\text{Hk14})_2$  determined in this work is consistent with previously published data determined using a fluorescently labeled hook peptide of the same sequence<sup>27</sup>. A comparison between these constants and the length of the peptide presented in Fig. 3 revealed two ranges where relatively small changes in the peptide length resulted in a substantial difference in complex stability. The largest increase in the stabilization of dimeric species,  $\sim 4.3$  orders of magnitude, was observed

between Hk5 and Hk14, corresponding to the  $\beta$ -hairpin structure, whereas an additional increase in affinity, by  $\sim 1.5$  orders of magnitude, was observed between Hk27 and Hk45, associated with an increase in the formation of the helical and coiled-coil structures observed in these peptide complexes (Fig. 2b). Similar dependence of the stability of zinc finger-like complexes on the length of peptide chain has been observed in previous studies, although in those works electrostatic or hydrophobic head-to-head or head-to-tail interactions were indicated as the reason for elevated stability<sup>25,28,29</sup>.

**Formation of the hydrogen bond network induced by  $\text{Zn}^{2+}$  binding.** Because the largest increase in the stabilization of dimeric zinc hook complexes occurred between the Hk5 and Hk14, we attempted to identify the structural features of these peptides and  $\text{Zn}^{2+}$  complexes. To this end, we examined the Hk4-Hk14 peptides in both metal-free and  $\text{Zn}^{2+}$  complexes using NMR and HDX MS. The 2D TOCSY, ROESY,  $^1\text{H}/^{13}\text{C}$  HSQC and  $^1\text{H}/^{15}\text{N}$  HSQC analyses facilitated the nearly complete resonance assignment of the  $^1\text{H}$ ,  $^{13}\text{C}$ , and  $^{15}\text{N}$  chemical shifts for the  $\text{Zn}^{2+}$  complexes of the studied zinc hook peptides (Table S2, Table S3). Analysis of the 2D ROESY spectrum of the  $\text{Zn}(\text{Hk14})_2$  complex generated 131 through-space contacts between proton pairs. A comparison of the experimental spectrum for  $\text{Zn}(\text{Hk14})_2$  with a simulated spectrum for the corresponding fragment of the crystal structure (PDB ID: IL8D<sup>6</sup>) revealed no differences in the central part of the  $\beta$ -hairpin, while terminal residues displayed higher mobility (and fewer ROE contacts) in the peptide compared to their counterparts in the crystal structure. These data showed that the structure of the  $\text{Zn}(\text{Hk14})_2$  complex observed in the solution is similar to the conformation of the corresponding fragment of the hook crystal structure. Using NMR, we measured the exchange rates and



temperature coefficients of amide protons in the zinc complexes of Hk6, Hk8, Hk10 and Hk14. Amide proton exchange is a powerful method to probe structural and dynamic properties at residue-level resolution, facilitating the distinction of solvent-exposed amides and amides within structured regions of the protein protected against exchange. The amide proton solvent accessibility is expressed as a protection factor representing the exchange rate expected for amide protons in unfolded peptides under a given condition, divided by the observed rate of exchange (Table S4). It can be safely assumed that amide protons cannot be buried deeply in the protein structure of a short peptide; therefore a low temperature coefficient and high protection factor of amide proton are indicative of involvement in the formation of a hydrogen bond<sup>30,31</sup>. The NMR results showed that amides of Cys444, Cys447, Gly448 and Arg449 are highly protected from the solvent in Zn(Hk14)<sub>2</sub> (Table S5), but no amides are protected from solvent exchange for the metal-free Hk14, evidenced by the 1D <sup>1</sup>H NMR spectra (Fig. S5). The same pattern of protected amides was also observed for Zn(Hk10)<sub>2</sub> but was less manifested in the Zn(Hk8)<sub>2</sub> and Zn(Hk6)<sub>2</sub>. The structural mapping of these highly protected amides supports a network of N-H...O and N-H...S hydrogen bonds within the  $\beta$ -hairpin formed upon Zn<sup>2+</sup> complexation (Fig. 4). These data showed that Zn<sup>2+</sup> binding nucleates folding of the  $\beta$ -hairpin of the hook domain. This mechanism was also supported by the results of HDX MS of Zn<sup>2+</sup> complexes of Hk4-Hk14, which showed three protected amide protons in the Zn<sup>2+</sup> complexes of Hk10-Hk14: one amide proton was identified within Cys444, while another two amide protons were detected in the Gly448-Leu451 fragment and two protected protons were observed in the Hk6 complex, localized on Cys444 and either Gly448 or Arg449 (see *Supplementary information* for details, Figs S6-S10).

**Impact of  $\beta$ -hairpin formation on stability of the hooked structure.** The increasing stability of the zinc hook complex with peptide length observed in Hk5-Hk14 supports the idea that the observed increase in stability is associated with the formation of the  $\beta$ -hairpin. Therefore we decided to investigate the isolated effect of the formation of the  $\beta$ -hairpin on the stability of the  $\text{Zn}(\text{Hk})_2$  complexes. Based on previous results showing that  $\beta$ -hairpin folding begins with turn formation and propagates toward the tail<sup>32,33</sup> and the metal-mediated folding of  $\beta\beta\alpha$  zinc fingers is initiated with the formation of the turn that nucleates the  $\beta$ -hairpin<sup>34</sup>, we hypothesized that the disruption of the turn region will prevent formation of the  $\beta$ -hairpin structure<sup>35,36</sup>. To this end, the amide of Gly448 was replaced with an ester linkage, resulting in a loss of the hydrogen-bond donor of the  $\beta$ -turn in *depsiHk* peptides (Fig. 3). The resulting *depsiHk8* and *depsiHk14* peptides exhibited significantly smaller changes in molar ellipticity at 222 nm compared with Hk8 or Hk14 upon  $\text{Zn}^{2+}$  binding (Fig. S11). The stability of  $\text{Zn}(\text{depsiHk8})_2$  and  $\text{Zn}(\text{depsiHk14})_2$  was significantly decreased by  $\sim 2.5$  and  $\sim 3$  orders of magnitude in the  $K_{12}$  values, respectively, compared with the parent  $\text{Zn}(\text{Hk8})_2$  and  $\text{Zn}(\text{Hk14})_2$  complexes (Fig. 3, Table 1 and S6). We combined an amide-to-ester substitution with the previously described alanine substitutions of conserved residues of the hydrophobic core (Hk14VA, Hk14LA, and Hk14VALA)<sup>27</sup>. The combined mutations of hydrophobic residues and amide-to-ester substitutions in the 14-amino acid fragment (*depsiHk14LA* and *depsiHk14VALA*) decreased the stability of the  $\text{Zn}(\text{Hk})_2$  species to the level of the  $\text{Zn}^{2+}$  complex formed by Hk4 peptide. These data indicate the additive effect of amide-to-ester substitutions and mutations of hydrophobic residues on the stability of the zinc hook complex, showing that hydrophobic interactions and backbone hydrogen bonding are the major structural factors governing the high stability of the complex (Fig. 3, Table 1 and Table S6).

Furthermore,  $\beta$ -turn-favoring Pro445 and hydrophobic Val446 in the spacer between two  $\text{Zn}^{2+}$  binding cysteines showed stabilizing effects, even in the context of minimal Hk4, as seen in Hk4VA, Hk4PA, and Hk4PAVA (Fig. 3, Table 1 and Table S6).

**Influence of the conserved hydrophobic core on formation of the coiled-coil.** We examined the effect of conserved hydrophobic residues (Val446 and Leu451) on the conformation and stability of the 45-mer zinc hook model by constructing peptides with either single or double substitutions to alanine (Hk45VA, Hk45LA and Hk45VALA). Compared with the Hk45 peptide, Hk45VA and Hk45LA showed two- and ten-fold less pronounced structural changes, respectively, upon the formation of the corresponding  $\text{Zn}(\text{Hk})_2$  complexes, evidenced by changes in the molar ellipticity at 222 nm (Fig. S12). The double-mutated Hk45VALA showed even smaller changes in molar ellipticity upon the formation of  $\text{Zn}(\text{Hk})_2$ , indicating a marginal degree of metal-coupled folding (Fig. S12). Mutations of Val446 and Leu451 significantly affected the stability of the  $\text{Zn}(\text{Hk})_2$  complex (Fig. 3, Table 1). The 2.85 order of magnitude decrease in stability, in terms of the  $K_{12}$  values between Hk45 and Hk45VALA, was larger than that between Hk14 and Hk14VALA (1.79 orders of magnitude)<sup>27</sup>. These data indicate that the interface formed by the Val446 and Leu451 residues of the  $\beta$ -hairpin and the hydrophobic residues of the coiled-coil, observed in the crystal structure, is essential for metal-coupled folding and the stable assembly of the hook domain.

**Acid-base properties of cysteine thiols in CXXC motif.** The acidity of the  $\text{Zn}^{2+}$  coordinating cysteines has a critical impact on the metal-binding properties as thiolate anions act as ligands to form complexes with transition metals<sup>37</sup>. The  $\text{p}K_{\text{a}}$  of a cysteine (here  $\text{p}K_{\text{a}}^{\text{SH}}$ ) can be strongly influenced by its protein microenvironment<sup>38,39</sup>. Thus, the deprotonation of

cysteine thiols was spectrophotometrically measured at 220 nm using metal-free forms of the Hk peptides (Fig. 5)<sup>15,40,41</sup>. The obtained  $pK_{a1}^{SH}$  and  $pK_{a2}^{SH}$  values were consistent with the values obtained in potentiometric titrations (Table S7, Fig. S13). Interestingly, a significant increase in the acidity of one cysteine thiol ( $pK_{a1}^{SH} = 8.2 - 7.5$ ) was observed as the peptide chain length increased from Hk4 to Hk10, whereas the  $pK_{a2}^{SH}$  of a second cysteine thiol remained constant at  $\sim 9.2$ . Further increases in the domain length up to Hk45 did not result in additional increases in thiol acidity (Table S7). These data showed that the 10-amino acid-long central region of the hook hairpin (Hk10) in the metal-free form captures the essential structural features necessary to accommodate the significantly perturbed  $pK_a$  values of the thiol group of one of the cysteine residues in the hook CXXC motif, thereby reducing the enthalpic cost of thiol deprotonation associated with  $Zn^{2+}$  complex formation (see below). Interestingly, the CXXC motif in the active site of certain enzymes, such as the thiol-disulfide oxidoreductases of the thioredoxin superfamily, is characterized by similarly decreased  $pK_a^{SH}$  of the N-terminal cysteine compared with the second cysteine.<sup>39,42</sup> We postulate that the CXXC motif in hook peptides might adopt a similar conformation, suggesting that the reduced  $pK_{a1}^{SH}$  value of  $\sim 7.5$  corresponds to Cys444. It has been proposed that  $Zn^{2+}$  assisted deprotonation of all the cysteines during the folding of  $Zn(Cys)_4$  zinc-finger cores and that the ionized cysteine core is stabilized by the interactions with the protein-derived structural elements<sup>25,43</sup>. The formation of N-H $\cdots$ S hydrogen bonds, which we also observed in the zinc hook structure, has been proposed to be an important factor for such stabilization<sup>44</sup>. Such stabilization of the  $Zn(Cys)_4$  core by the protein-derived structural elements is expected to result in both stabilization of the  $Zn^{2+}$  complex and increase in the apparent acidity of the peptide thiols in the presence of  $Zn^{2+}$ . To test this prediction, we performed spectrophotometric pH titrations of zinc hook peptides at a 1:2  $Zn^{2+}$ -to-peptide molar ratio. The results of these titrations were used to calculate the apparent average value of

peptide thiols' deprotonation in the presence of  $\text{Zn}^{2+}$  ions ( $\text{pK}_a'$ ), which reflects the competition between protons and  $\text{Zn}^{2+}$  for binding to cysteine thiolates<sup>15,28</sup>. The obtained  $\text{pK}_a'$  values (from 6.25 for Hk4 to 4.70 for Hk45, Table S8) and apparent formation constants ( $\log K_{12}$ ) of the zinc complexes are highly correlated (Table 1, Fig. S14), which is consistent with the notion of the  $\text{Zn}^{2+}$ -assisted deprotonation the cysteines, and stabilization of the  $\text{Zn}(\text{Cys})_4$  core by the protein-derived structural elements. The  $\text{pK}_a'$  values of 5.1-5.6 were reported for the  $\text{Zn}^{2+}$ -induced Cys deprotonation in the case of  $(\text{Cys})_4$  and  $(\text{Cys})_3(\text{His})$  zinc fingers<sup>14,29,37,45</sup>. Recently, the  $\text{pK}_a'$  value of cysteines between 4.2 and 5.1 was reported for the artificial  $(\text{Cys})_4$  treble-clef zinc finger  $\text{L}_{\text{TC}}$ , but the exact deprotonation value was not calculated<sup>25</sup>. It should be noted that the  $\text{pK}_a'$  values obtained in different studies cannot be directly compared, because  $\text{pK}_a'$  values are specific for particular conditions (i.e. concentrations and  $\text{Zn}^{2+}$ -to-peptide ratios).

**Thermodynamics of zinc hook domain formation.** In order to study the thermodynamics of  $\text{Zn}^{2+}$  binding to Hk4-Hk45 peptides in detail, we used ITC to determine the enthalpies of formation of the corresponding  $\text{Zn}(\text{Hk})_2$  complexes (Fig. 6 and Fig. S15). The resulting experimental enthalpies ( $\Delta H_{\text{ITC}}$ ) of the formation of  $\text{Zn}(\text{Hk})_2$  complexes are presented in Table 2, and the complete fit values are presented in Table S9. It should be noted that same enthalpies of the formation of  $\text{Zn}(\text{Hk})_2$  were obtained by both titrations of  $\text{Zn}^{2+}$  ions into peptide and titrations of peptide into  $\text{Zn}^{2+}$  (Figs S16 and S17). Because  $\text{Zn}^{2+}$  complexation by cysteine-containing peptides is accompanied by thiol deprotonation, the experimental enthalpy ( $\Delta H_{\text{ITC}}$ ) must be corrected for the heat of ionization of the buffer<sup>46,47</sup>, as indicated in equation (4):

$$\Delta H_{\text{ITC}} = \Delta H^\circ + n_{\text{H}} \Delta H^\circ_{\text{buff}} \quad (4)$$

Where  $\Delta H^\circ$  is the buffer independent intrinsic reaction enthalpy,  $n_H$  is the number of protons released over the course of the reaction, and  $\Delta H^\circ_{\text{buff}}$  is the buffer-specific heat of ionization (-5.02 kcal/mol for HEPES)<sup>48</sup>. The number of released protons upon  $\text{Zn}^{2+}$  binding is the number of protons associated with cysteine thiols of two metal-free peptide molecules at pH 7.4, calculated based on the  $\text{p}K_{\text{a}1}^{\text{SH}}$  and  $\text{p}K_{\text{a}2}^{\text{SH}}$  values (Table S7)<sup>49</sup>. The resulting buffer-independent intrinsic reaction enthalpy,  $\Delta H^\circ$ , can be further dissected into two components: the enthalpy of  $\text{Zn}^{2+}$  binding to the peptide and the associated structural changes ( $\Delta H^\circ_{\text{Zn-pep}}$ ) and enthalpy of deprotonation of cysteine thiols associated with  $\text{Zn}^{2+}$  binding ( $n_H \Delta H^\circ_{\text{CysH}}$ )<sup>14</sup>, as expressed in equation (5):

$$\Delta H^\circ = \Delta H^\circ_{\text{Zn-pep}} + n_H \Delta H^\circ_{\text{CysH}} \quad (5)$$

where  $n_H$  is the number of protons released over the course of the reaction (see above), and  $\Delta H^\circ_{\text{CysH}}$  is the heat of Cys deprotonation. The average value of the last parameter is +8.5 kcal/mol<sup>50</sup>, and it is assumed that it is constant in the studied series of peptides. In addition to these factors, the entropic component of the reaction ( $-\text{T}\Delta S^\circ$ ) was calculated based on Gibbs' law, as shown in equation (6):

$$\Delta G^\circ = \Delta H^\circ - \text{T}\Delta S^\circ \quad (6)$$

This equation uses the  $\Delta G^\circ$  values that were calculated based on the apparent formation constants ( $K_{12}$ ) of the hook complexes from Table 1 as indicated in equation (7):

$$\Delta G^\circ = -\text{RTln}K_{12} \quad (7)$$

A comparison of the  $\Delta G^\circ$ ,  $\Delta H^\circ$ , and  $-\text{T}\Delta S^\circ$  listed in Table 2 revealed that the formation of the zinc hook complexes is largely entropically driven. The entropically driven binding of  $\text{Zn}^{2+}$  has been demonstrated to be likely an intrinsic property of  $\text{Zn}(\text{Cys})_4$  coordination spheres in proteins<sup>14</sup>, whereas  $\text{Zn}^{2+}$  binding to the sites containing

histidine ligands are more entropically (and less enthalpically) driven as the number of histidine ligands increases<sup>49,51</sup>. The increase in stability of the hook complexes with the peptide length, reflected as the decrease of the  $\Delta G^\circ$  values, results primarily from the increase in favorable enthalpy ( $\Delta H^\circ$ ). The observed dependence of the enthalpy on the domain fragment length is partially compensated by the unfavorable change in entropy ( $-T\Delta S^\circ$ ), demonstrating entropy-enthalpy compensation (EEC)<sup>52</sup>. As evident in Figure 7, EEC was only observed in the series of peptides where increasing peptide length paralleled an increasing tendency for secondary structure formation resulting from  $\text{Zn}^{2+}$  binding (Fig. 2b):  $\beta$ -hairpin in Hk4-Hk8 and coiled-coil in Hk31-Hk45. For these peptide series, the enthalpy of binding becomes more favorable, and the entropy becomes more unfavorable, as the length of the structured fragment increases, suggesting that the favorable increase in enthalpy is closely correlated with the interactions resulting from metal-mediated folding; however, this increase is partially compensated by the unfavorable loss in entropy resulting from the formation of ordered structures and the association of the subunits. Next, we analyzed in detail the thermodynamic contributions from the structural components present in the hook domain to the stability of the hook complex. The minimal hook peptide Hk4 can serve as a reference for the  $\text{Zn}(\text{Cys}_2)_2$  zinc hook coordination motif in the absence of other protein-derived structural elements (Fig. 7). The  $\Delta H^\circ_{\text{Zn-pep}}$  value for  $\text{Zn}(\text{Hk4})_2$  (-25 kcal/mol) was identical to that obtained for the Gly-rich peptide previously used to model the  $\text{Zn}(\text{Cys})_4$  coordination motif of structural zinc sites, such as those found in zinc fingers, in the absence of protein-derived structural elements, indicating that the enthalpy of  $\text{Zn}^{2+}$ -S bond formation is similar in these two models<sup>14,37</sup>. In order to obtain information on the thermodynamic contributions from the structural components present in the hook domain, we compared the thermodynamic parameters

( $\Delta G^\circ$ ,  $\Delta H^\circ$ ,  $-T\Delta S^\circ$ ,  $\Delta H^\circ_{\text{Zn-pep}}$ ,  $n_{\text{H}}\Delta H^\circ_{\text{CysH}}$ ) for the formation of  $\text{Zn(Hk)}_2$  complexes of the peptides that encompass particular structural fragments (Table 2). Subtraction of the thermodynamic parameters ( $\Delta G^\circ$ ,  $\Delta H^\circ$ ,  $-T\Delta S^\circ$ ,  $\Delta H^\circ_{\text{Zn-pep}}$ ,  $n_{\text{H}}\Delta H^\circ_{\text{CysH}}$ ) for  $\text{Zn(Hk4)}_2$  from the corresponding values for  $\text{Zn(Hk14)}_2$  and subtraction of the thermodynamic parameters for  $\text{Zn(Hk14)}_2$  from the corresponding values for  $\text{Zn(Hk45)}_2$  gave the information of the net thermodynamic effect of the protein-derived structural elements in the  $\beta$ -hairpin ( $\beta$ ) and coiled-coil (cc) domain fragments respectively (Fig. 8). These analyses show that the formation of the  $\text{Zn(Cys)}_2$  coordination motif (minimally as  $\text{Zn(Hk4)}_2$ ) provides -20.4 kcal/mol of Gibbs free energy, while protein-derived structural elements in the  $\beta$ -hairpin and the coiled-coil regions provide an additional -5.8 kcal/mol and -2.1 kcal/mol of free energy, respectively. The change in free energy emerges from favorable change in enthalpy ( $\Delta H^\circ$ ) of -8.9 kcal/mol and -4.5 kcal/mol and an unfavorable entropic cost ( $-T\Delta S^\circ$ ) of 3.1 kcal/mol and 2.4 kcal/mol for  $\beta$ -hairpin and coiled-coil regions respectively. The analysis of the enthalpic component ( $\Delta H^\circ$ ) according to equation (5) shows that the  $\beta$ -hairpin formation contributes -4.3 kcal/mol to the enthalpy due to the favorable change in enthalpy of  $\text{Zn}^{2+}$  binding and the associated structural changes ( $\Delta\Delta H^\circ_{\text{Zn-pep}}$ ) and -4.5 kcal/mol due to the decrease in the energetic cost of cysteine deprotonation ( $\Delta(n_{\text{H}}\Delta H^\circ_{\text{CysH}})$ ). Correspondingly, the coiled-coil fragment contributes -4.3 kcal/mol to the favorable change in enthalpy of  $\text{Zn}^{2+}$  binding and the associated structural changes ( $\Delta\Delta H^\circ_{\text{Zn-pep}}$ ), providing only a negligible decrease in the energetic cost of cysteine deprotonation ( $\Delta(n_{\text{H}}\Delta H^\circ_{\text{CysH}})$ ) (Fig. 8). This energetic effect is directly associated with the increased acidity of one of the thiols in the metal-free form, occurring almost entirely in the  $\beta$ -hairpin-forming fragment (Fig. 5). These data show that the contribution to the stability of the  $\beta$ -hairpin-forming fragment is evident in both metal-free and  $\text{Zn}^{2+}$ -bound forms. In the



metal-free form, an increase of the acidity of one of the thiols decreases the unfavorable enthalpy of deprotonation, whereas in the  $\text{Zn}^{2+}$  complex, a network of hydrogen bonds and other interactions is formed, increasing the favorable enthalpy of the reaction.

**Comparison with the stability of other zinc binding domains.** Currently, very limited data on metal ion affinity for intermolecular  $\text{Zn}^{2+}$  binding domains exist. One example of a relatively well-studied domain is the intermolecular zinc binding site formed by CD4/CD8 $\alpha$  co-receptors and Lck kinase. The apparent formation constants ( $\log K$ ) were reported as 6.4 for CD4-Lck and 6.05 for CD8 $\alpha$ -Lck<sup>53</sup>. However, these values were measured in the presence of excess  $\text{Zn}^{2+}$ , and thus the metal concentration factor was neglected in the calculation of the stability constant. In the following study, the authors considered both the metal and peptide concentrations and measured the apparent stability constants of  $\text{Co}^{2+}$ -mediated dimers formed by minimal peptides from CD4, CD8 $\alpha$  and Lck reporting values, which expressed as apparent formation constants ( $\log K$ ) are 7.8 for the CD4- $\text{Co}^{2+}$ -Lck complex and 8.1 for the CD8 $\alpha$ - $\text{Co}^{2+}$ -Lck complex<sup>54</sup>.

In contrast to intermolecular domains, intramolecular zinc binding domains, such as zinc fingers, have been studied in considerable detail in terms of their metal-binding affinity. The typical values of  $\log K$  reported for natural zinc binding fingers ranges from 10 to 14<sup>1,14,15,29,55,56</sup>. The stability values reported for intramolecular ( $\text{ZnL}$ ) complexes are not directly comparable with the constants of dimeric complexes ( $\text{ZnL}_2$ ) presented here (Table 1) because of the different definitions of the equilibrium constants. An indirect way to compare these conditional equilibrium constants is to analyze relative complex formation in a setting where both peptides compete for  $\text{Zn}^{2+}$  using e.g. Hyperquad Simulation and Speciation software<sup>57</sup>. Our analysis revealed that the zinc hook domain would thermodynamically outcompete the

typical or even the strongest natural zinc finger domains (Fig. S18). These results show that zinc hook and other similar intermolecular  $\text{Zn}^{2+}$  binding sites in proteins can form metal-mediated assemblies at very low concentrations of subunits under physiologically buffered concentrations of  $\text{Zn}^{2+}$  ions<sup>58</sup>.

**Conclusions.** The present study provides a detailed analysis of the determinants of intermolecular Rad50 zinc hook domain stabilization using model peptides of gradually increasing lengths and mutational variants. Although we anticipated that the  $\text{Zn}^{2+}$  affinity would increase with hook peptide length, thereby increasing the total number of intra- and inter-molecular interactions, most of the stabilization effect reflected a small portion of the Rad50 hook domain surrounding the  $\text{Zn}^{2+}$  binding motif. The significant stabilization of the hook complex resulted from the favorable increase in the enthalpy of  $\text{Zn}^{2+}$  complexation. This reflects a reduction of the unfavorable enthalpy of Cys thiol deprotonation ascribed to the  $\beta$ -hairpin forming fragment and the favorable enthalpy of interactions in the  $\beta$ -hairpin and coiled-coil structures of the domain, which are formed upon metal binding. We postulated that this metal-coupled folding dictates molecular recognition and the specificity and stability of the interaction and is responsible for the long-range allostery observed between the hook and globular domains of Rad50 in recent studies<sup>11,13,23</sup>.

## Methods

**Peptide synthesis.** Zinc hook peptides (Hk) were synthesized via solid-phase synthesis (SPPS) using an Fmoc strategy. All peptides were N-terminally acetylated. Amide-to-ester backbone bond-substituted peptide analogs (*depsi*Hk) were synthesized according to Jemth and co-workers, followed by fragment condensation<sup>35</sup>.

**Protonation and  $\text{Zn}^{2+}$  stability constants.** The protonation and  $\text{Zn}^{2+}$  stability constants of Hk4-Hk14 zinc hook peptides and depsipeptides were determined in potentiometric titrations. The apparent formation constants of  $\text{Zn}^{2+}$  complexes with zinc hook peptides (Hk23-45, Hk45VA, Hk45LA, and ZnHK45VALA) were determined in the spectropolarimetric titrations with  $\text{Zn}^{2+}$  in the presence of HEDTA, EDTA and TPEN chelators<sup>15,27</sup>. The  $\text{p}K_{a1}^{\text{SH}}$ ,  $\text{p}K_{a2}^{\text{SH}}$  dissociation constants of the cysteine thiols of Hk4-Hk45 were determined spectrophotometrically as described previously<sup>41</sup>. Similarly, the dissociation constants of thiols in the presence of  $\text{Zn}^{2+}$  ( $\text{p}K_a'$ ) were determined by the pH titration of Hk4-Hk14 and Hk45 in the presence of 0.495 eq. of  $\text{Zn}^{2+}$ .

**NMR studies.** NMR measurements of 5 mM  $\text{Zn}(\text{Hk6})_2$ ,  $\text{Zn}(\text{Hk10})_2$ ,  $\text{Zn}(\text{Hk12})_2$  and  $\text{Zn}(\text{Hk14})_2$  and metal-free Hk14 were performed in degassed 10%  $\text{D}_2\text{O}$  in  $\text{H}_2\text{O}$  (pH 7.4) on a DDR2 Agilent 600 MHz spectrometer equipped with a Penta probe.

**Hydrogen-deuterium exchange mass spectrometry (HDX MS).** The experiments were conducted in the exchange-in mode, e.g., hydrogen into deuterium. For each deuterated sample, isotopic profiles of 1:1 and 1:2 metal-to-peptide complexes were compared with the theoretical isotopic profile of fully deuterated species, and the signals corresponding to complexes with various numbers of protected protons were analyzed. Subsequently, the fragmentation ions were analyzed in the same manner.

**Isothermal titration calorimetry (ITC).** The binding of  $\text{Zn}^{2+}$  to Hk peptides was monitored using ITC at 25°C. All experiments were performed in HEPES buffer ( $I = 0.1$  M from NaCl) at pH 7.4 under an argon atmosphere. The Hk peptide (titrant) concentration was 1.3 mM, whereas the metal (titrate) concentration was 50  $\mu\text{M}$ . The titration data were fitted to a

binding model accounting for the formation of ZnHk and Zn(Hk)<sub>2</sub> complexes during the course of titration.

For a more detailed description of the experimental methods of peptide synthesis and zinc hook complex characterization (NMR, ITC, HDX MS, UV-vis, and circular dichroism spectroscopy), see Supplementary information.

## **Acknowledgements**

This research was financially supported by grants from the National Science Centre (NCN) under OPUS grant no. 2014/13/B/NZ1/00935 and partly by a Ministry of Science and Higher Education grant (no. DI2011 031341). The instrumentation was financially supported by the Polish Foundation for Science under Focus projects F1/2010/P/2013 (A.K.\*) and FG1/2010 (A.K.\*) and the European Union from the European Regional Development Fund under the Operational Program Innovative Economy, 2007-2013. The authors would like to thank Prof. Wojciech Bal, Dr. Tomasz Frączyk, and Dr. Wojciech Jachymek for providing the ITC and ESI-MS instrumentation and for their kind discussions. T.K., M.N., D.W., W.K. and A.K.\* would like to thank the Foundation for Polish Science for support by the FOCUS and TEAM programs.

## **Authors' contributions**

T.K., M.N., D.W., A.K., and A.K.\* performed the experiments and analyzed the data. T. K. and A.K.\* designed the research in consultation with M.N., A.E., and W.K. T.K. and A.K.\* wrote the manuscript. The authors declare no competing financial interests.

## Additional information

The authors declare no competing financial interest.

## References

1. Kochańczyk, T., Drozd, A. & Krężel, A. Relationship between the architecture of zinc coordination and zinc binding affinity in proteins - insights into zinc regulation. *Metallomics* **7**, 244–257 (2015).
2. Maret, W. & Li, Y. Coordination dynamics of zinc in proteins. *Chem. Rev.* **109**, 4682–4707 (2009).
3. Lepore, B. W., Ruzicka, F. J., Frey, P. A. & Ringe, D. The x-ray crystal structure of lysine-2,3-aminomutase from *Clostridium subterminale*. *Proc. Natl. Acad. Sci. U.S.A.* **102**, 13819–13824 (2005).
4. Schmalen, I. *et al.* Interaction of circadian clock proteins CRY1 and PER2 is modulated by zinc binding and disulfide bond formation. *Cell* **157**, 1203–1215 (2014).
5. Kim, P. W., Sun, Z. Y., Blacklow, S. C., Wagner, G. & Eck, M. J. A zinc clasp structure tethers Lck to T cell coreceptors CD4 and CD8. *Science* **301**, 1725–1728 (2003).
6. Hopfner, K. P. *et al.* The Rad50 zinc-hook is a structure joining Mre11 complexes in DNA recombination and repair. *Nature* **418**, 562–566 (2002).
7. Ciszak, E. & Smith, G. D. Crystallographic evidence for dual coordination around zinc in the T3R3 human insulin hexamer. *Biochemistry* **33**, 1512–1517 (1994).
8. Bixby, K. A. *et al.* Zn<sup>2+</sup>-binding and molecular determinants of tetramerization in voltage-gated K<sup>+</sup> channels. *Nat. Struct. Biol.* **6**, 38–43 (1999).

9. Stracker, T. H. & Petrini, J. H. The MRE11 complex: starting from the ends. *Nat. Rev. Mol. Cell. Biol.* **12**, 90–103 (2011).
10. Herdendorf, T. J., Albrecht, D. W., Benkovic, S. J. & Nelson, S. W. Biochemical characterization of bacteriophage T4 Mre11-Rad50 complex. *J. Biol. Chem.* **286**, 2382–2392 (2010).
11. Roset, R. *et al.* The Rad50 hook domain regulates DNA damage signaling and tumorigenesis. *Genes Dev.* **28**, 451–462 (2014).
12. Barfoot, T. *et al.* Functional analysis of the bacteriophage T4 Rad50 homolog (gp46) coiled-coil domain. *J. Biol. Chem.* **290**, 23905–23915 (2015).
13. Hohl, M. *et al.* Interdependence of the Rad50 hook and globular domain functions. *Mol. Cell* **57**, 479–491 (2015).
14. Reddi, A. R., Guzman, T. R., Breece, R. M., Tierney, D. L. & Gibney, B. R. Deducing the energetic cost of protein folding in zinc finger proteins using designed metallopeptides. *J. Am. Chem. Soc.* **129**, 12815–12827 (2007).
15. Miłoch, A. & Krężel, A. Metal binding properties of the zinc finger metallome--insights into variations in stability. *Metallomics* **6**, 2015–2024 (2014).
16. Song, W. J., Sontz, P. A., Ambroggio, X. & Tezcan, F. A. Metals in protein-protein interfaces. *Annu. Rev. Biophys.* **43**, 409–431 (2014).
17. Nooren, I. M. & Thornton J. M. Diversity of protein-protein interactions. *EMBO J.* **22**, 3486–3492 (2003).
18. Jones, S. & Thornton, J. M. Principles of protein-protein interactions. *Proc. Natl. Acad. Sci. U.S.A.* **93**, 13–20 (1996).
19. Acuner Ozbabacan, S. E., Engin, H. B., Gursoy A. & Keskin, O. Transient protein-protein interactions. *Protein. Eng. Des. Sel.* **24**, 635–648 (2011).

20. Krissinel, E. & Kenrick, E. Inference of macromolecular assemblies from crystalline state. *J. Mol. Biol.* **372**, 774–797 (2007).
21. Krężel, A., Latajka, R., Bujacz, G. & Bal, W. Coordination properties of tris(2-carboxyethyl)phosphine, a newly introduced thiol reductant, and its oxide. *Inorg. Chem.* **42**, 1994–2003 (2003).
22. Perczel, A., Hollosi, M., Foxman, B. M. & Fasman, G. D. Conformational analysis of pseudocyclic hexapeptides based on quantitative circular dichroism (CD), NOE, and X-ray data. The pure CD spectra of type I and type II beta-turns. *J. Am. Chem. Soc.* **113**, 9772–9784 (1991).
23. Graddis, T. J., Myszkowski, D. G. & Chaiken, I. M. Controlled formation of model homo- and heterodimer coiled coil polypeptides. *Biochemistry* **32**, 12664–12671 (1993).
24. Hohl, M. *et al.* The Rad50 coiled-coil domain is indispensable for Mre11 complex functions. *Nat. Struct. Mol. Biol.* **18**, 1124–1131 (2011).
25. Sénèque, O., Bonnet, E., Joumas, F. L. & Latour, J. M. Cooperative metal binding and helical folding in model peptide of treble-clef zinc fingers. *Chem. Eur. J.* **15**, 4798–4810 (2009).
26. Krężel, A. & Maret, W. The biological inorganic chemistry of zinc ions. *Arch. Biochem. Biophys.* doi: 10.1016/j.abb.2016.04.010 (2016).
27. Kochańczyk, T., Jakimowicz, P. & Krężel, A. Femtomolar Zn(II) affinity of minimal zinc hook peptides - A promising small tag for protein engineering. *Chem. Commun. (Camb)* **49**, 1312–1314 (2013).
28. Sikorska, M., Krężel, A. & Otlewski, J. Femtomolar Zn<sup>2+</sup> affinity of LIM domain of PDLIM1 protein uncovers crucial contribution of protein-protein interactions to protein stability. *J. Inorg. Biochem.* **115**, 28–35 (2012).

29. Bombarda, E. *et al.* Determination of the  $pK_a$  of the four  $Zn^{2+}$ -coordinating residues of the distal finger motif of the HIV-1 nucleocapsid protein: consequences on the binding of  $Zn^{2+}$ . *J. Mol. Biol.* **310**, 659–672 (2001).
30. Wójcik, J., Ruszczyńska, K., Zhukov, I. & Ejchart, A. NMR measurements of proton exchange between solvent and peptides and proteins. *Acta Biochim. Polon.* **46**, 651–663 (1998).
31. Bai, Y., Milne, J. S., Mayne, L. & Englander, S. W. Primary structure effects on peptide group hydrogen exchange. *Proteins Structure Funct. Bioinf.* **17**, 75–86 (1993).
32. Munoz, V., Thompson, P. A., Hofrichter, J. & Eaton, W. A. Folding dynamics and mechanism of  $\beta$ -hairpin formation. *Nature* **390**, 196–199 (1997).
33. Du, D., Zhu, Y., Huang, C.-Y. & Gai, F. Understanding the key factors that control the rate of  $\beta$ -hairpin folding. *Proc. Natl. Acad. Sci. U.S.A.* **101**, 15915–15920 (2004).
34. Li, W., Zhang, J., Wang, J. & Wang, W. Metal-coupled folding of Cys2His2 zinc-finger. *J. Am. Chem. Soc.* **130**, 892–900 (2008).
35. Eildal, J. N. *et al.* Probing the role of backbone hydrogen bonds in protein-peptide interactions by amide-to-ester mutations. *J. Am. Chem. Soc.* **135**, 12998–13007 (2013).
36. Deechongkit, S., Dawson, P. E. & Kelly, J. W. Toward assessing the position-dependent contributions of backbone hydrogen bonding to beta-sheet folding thermodynamics employing amide-to-ester perturbations. *J. Am. Chem. Soc.*, **126**, 16762–16771 (2004).
37. Reddi, A. R. & Gibney, B. R. Role of protons in the thermodynamic contribution of a Zn(II)-Cys4 site toward metalloprotein stability. *Biochemistry* **46**, 3745–3758 (2007).



38. Maynard, A. T. & Covell, D. G. Reactivity of zinc finger cores: analysis of protein packing and electrostatic screening. *J. Am. Chem. Soc.* **123**, 1047–1058 (2001).
39. Foloppe, N. & Nilsson, L. The glutaredoxin -C-P-Y-C- motif: influence of peripheral residues. *Structure* **12**, 289–300 (2004).
40. Krężel, A. & Maret, W. Dual nanomolar and picomolar Zn(II) binding properties of metallothionein. *J. Am. Chem. Soc.* **129**, 10911–10921 (2007).
41. Adamczyk, J., Bal, W. & Krężel, A. Coordination properties of dithiobutylamine (DTBA), a newly introduced protein disulfide reducing agent. *Inorg. Chem.* **54**, 596–606 (2015).
42. Van Laer, K., Oliveira, M., Wahni, K. & Messens, J. The concerted action of a positive charge and hydrogen bonds dynamically regulates the  $pK_a$  of the nucleophilic cysteine in the NrdH-redoxin family. *Protein Sci.* **23**, 238–242 (2013).
43. Dudev, T. & Lim, C. Factors governing the protonation state of cysteines in proteins: An ab initio/CDM study. *J. Am. Chem. Soc.* **124**, 6759–6766 (2002).
44. Maynard, A. T. & Covell, D. G. Reactivity of zinc finger cores: Analysis of protein packing and electrostatic screening. *J. Am. Chem. Soc.* **123**, 1047–1058 (2001).
45. Petros, A. K., Reddi, A. R., Kennedy, M. L., Hyslop, A. G. & Gibney, B. R. Femtomolar Zn(II) affinity in a peptide-based ligand designed to model thiolate-rich metalloprotein active sites. *Inorg. Chem.* **45**, 9941–9958 (2006).
46. Quinn, C. F., Carpenter, M. C., Croteau, M. L. & Wilcox, D. E. Isothermal Titration Calorimetry Measurements of Metal Ions Binding to Proteins. *Methods Enzymol.* **567**, 3–21, 2016
47. Johnson, R. A., Manley, O. M., Spuches, A. M. & Grosseohme, N. E. Dissecting ITC data of metal ions binding to ligands and proteins. *Biochim. Biophys. Acta* **1860**, 892–901 (2016).

48. Fukada, H. & Takahashi, K. Enthalpy and heat capacity changes for the proton dissociation of various buffer components in 0.1 M potassium chloride. *Proteins* **33**, 159–166 (1998).
49. Rich, A. M. *et al.* Thermodynamics of  $\text{Zn}^{2+}$  binding to Cys2His2 and Cys2HisCys zinc fingers and a Cys4 transcription factor site. *J. Am. Chem. Soc.* **134**, 10405–10418 (2012).
50. Wrathall, D., Izatt, R. & Christensen J., Thermodynamics of proton dissociation in aqueous solution. III. L-cysteine, S-methyl-L-cysteine, and mercaptoacetic acid. Determination of cysteine microconstants from calorimetric data. *J. Am. Chem. Soc.* **86**, 5712–5712 (1964).
51. Carpenter, M. C. & Wilcox, D. E. Thermodynamics of formation of the insulin hexamer: metal-stabilized proton-coupled assembly of quaternary structure. *Biochemistry* **53**, 1296–1301 (2014).
52. Liu, L. & Guo, Q. Isokinetic relationship, isoequilibrium relationship, and enthalpy-entropy compensation. *Chem. Rev.* **101**, 673–696 (2001).
53. Kim, P. W., Sun, Z. Y., Blacklow, S. C., Wagner, G. & Eck, M. J. A zinc clasp structure tethers Lck to T cell coreceptors CD4 and CD8. *Science* **301**, 1725–1728 (2003).
54. Davis, A. M. & Berg, J. M. Homodimerization and heterodimerization of minimal zinc(II)-binding-domain peptides of T-cell proteins CD4, CD8alpha, and Lck. *J. Am. Chem. Soc.* **131**, 11492–11497 (2009).
55. Posewitz, M. C. & Wilcox, D. E. Properties of the Sp1 zinc finger 3 peptide: coordination chemistry, redox reactions, and metal binding competition with metallothionein. *Chem. Res. Toxicol.* **8**, 1020–1028 (1995).

56. Imanishi, M. *et al.* Zn(II) Binding and DNA binding properties of ligand-substituted CXHH-type zinc finger proteins. *Biochemistry* **51**, 3342–3348 (2012).
57. Alderighi, L. *et al.* Hyperquad simulation and speciation (HySS): a utility program for the investigation of equilibria involving soluble and partially soluble species, *Coord. Chem. Rev.* **184** (1999) 311-318.
58. Krężel, A. & Maret, W. Zinc-buffering capacity of a eukaryotic cell at physiological pZn. *J. Biol. Inorg. Chem.* **11**, 1049–1062 (2006).

**Table 1.** Apparent formation constants ( $\log K_{12}$ ) of  $\text{Zn}(\text{Hk})_2$  complexes of zinc hook peptides and their mutants at pH 7.4,  $I = 0.1 \text{ M}$ ,  $25^\circ\text{C}$ . Values were calculated from cumulative protonation and stability constants determined potentiometrically (Tables S1 and S6) or determined spectropolarimetrically in competition experiments with zinc chelators (Fig. S4) and are not the results of ITC data fits.

Hook peptide	$\log K_{12}$	Mutant peptide	$\log K_{12}$
Hk4	$14.93 \pm 0.02$	Hk4VA	$14.41 \pm 0.02$
Hk5	$15.37 \pm 0.02$	Hk4PA	$14.19 \pm 0.02$
Hk6	$16.44 \pm 0.02$	Hk4PAVA	$13.96 \pm 0.02$
Hk8	$17.78 \pm 0.01$	Hk45LA	$19.06 \pm 0.08$
Hk10	$18.64 \pm 0.01$	Hk45VA	$19.78 \pm 0.08$
Hk12	$19.02 \pm 0.01$	Hk45LAVA	$17.89 \pm 0.08$
Hk14	$19.19 \pm 0.01$		
Hk23	$19.49 \pm 0.04$	Depsipeptide	$\log K_{12}$
Hk27	$19.77 \pm 0.04$	<i>depsiHk8</i>	$15.33 \pm 0.01$
Hk31	$20.47 \pm 0.09$	<i>depsiHk14</i>	$16.21 \pm 0.01$
Hk37	$20.69 \pm 0.06$	<i>depsiHk14LA</i>	$14.73 \pm 0.03$
Hk45	$20.74 \pm 0.06$	<i>depsiHk14VALA</i>	$13.97 \pm 0.02$

**Table 2.** Thermodynamic parameters of Zn(Hk)<sub>2</sub> complex formation at pH 7.4, *I* = 0.1 M, 25°C. The  $\Delta G^\circ$  values were calculated based on apparent formation constants  $K_{12}$  presented in Table 1 using equation (7).

Zinc hook peptide	$\Delta G^\circ$ (kcal/mol)	$\Delta H_{\text{ITC}}$ (kcal/mol)	$\Delta H^\circ$ (kcal/mol)	$\Delta H^\circ_{\text{Zn-pep}}$ (kcal/mol)	$n_{\text{H}}\Delta H^\circ_{\text{CysH}}$ (kcal/mol)	$-T\Delta S^\circ$ (kcal/mol)
Hk4	$-20.37 \pm 0.03$	$-12.1 \pm 0.03$	6.54	-25.08	31.62	-26.92
Hk5	$-20.97 \pm 0.03$	$-13.5 \pm 0.02$	4.75	-26.19	30.94	-25.73
Hk6	$-22.43 \pm 0.03$	$-15.5 \pm 0.02$	2.15	-27.77	29.92	-24.58
Hk8	$-24.26 \pm 0.01$	$-19.3 \pm 0.02$	-1.85	-31.43	29.58	-22.41
Hk10	$-25.43 \pm 0.01$	$-18.2 \pm 0.02$	-2.16	-29.36	27.2	-23.28
Hk12	$-25.95 \pm 0.01$	$-18.3 \pm 0.02$	-2.26	-29.46	27.2	-23.70
Hk14	$-26.18 \pm 0.01$	$-18.2 \pm 0.02$	-2.36	-29.22	26.86	-23.82
Hk23	$-26.59 \pm 0.05$	$-17.7 \pm 0.03$	-2.17	-28.52	26.35	-24.43
Hk27	$-26.97 \pm 0.05$	$-18.1 \pm 0.03$	-2.57	-28.92	26.35	-24.41
Hk31	$-27.9 \pm 0.01$	$-19.5 \pm 0.03$	-3.88	-30.40	26.52	-24.03
Hk37	$-28.22 \pm 0.08$	$-20.6 \pm 0.04$	-4.88	-31.57	26.69	-23.35
Hk45	$-28.29 \pm 0.08$	$-22.5 \pm 0.03$	-6.87	-33.39	26.52	-21.43

## Figure legends

**Figure 1.** Structure of the central fragment of the Rad50 protein from *P. furiosus*. a) Crystal structure of the 103-amino acid-long central fragment of Rad50, representing the zinc hook domain (PDB code: 1L8D)<sup>6</sup>; b) Sequence of the 45-amino acid Hk45 protein fragment used as a domain model in the present study. The dots indicate all residues forming the dimer interface, obtained using the PDBePISA web server. Val446 and Leu451 (green) residues were examined for the formation of hydrophobic interactions<sup>19</sup>. Cys444 and Cys447, which participate in Zn<sup>2+</sup> binding, are shown in yellow. The underlined residues participate in coiled-coil formation; c) Structural representation of the 45-amino acid-long fragment of the domain; d) Zinc hook peptides (Hk23-Hk45) used in the present study to probe the effect of coiled-coil formation; and e) Zinc hook peptides (Hk4-Hk14) used in the present study to probe the effect of  $\beta$ -hairpin formation.

**Figure 2.** CD spectra of zinc hook peptides recorded in 10 mM Tris-HCl buffer, 100 mM NaClO<sub>4</sub>, 100-200  $\mu$ M TCEP, pH 7.4. a) Titrations of Hk4, Hk8, Hk14, and Hk45 with Zn<sup>2+</sup> ions. The insets show the dependence of ellipticity at 220 nm on the Zn<sup>2+</sup>/Hk molar ratio. Red and blue colors refer to spectra of metal-free hook peptides and Zn(Hk)<sub>2</sub> complexes, respectively; b) Differential CD spectra of the zinc hook peptides obtained by the subtraction of spectra of free peptides from the spectra of Zn(Hk)<sub>2</sub> complexes.

**Figure 3.** Relationship between the apparent formation constants ( $\log K_{12}$ ) of the Zn(Hk)<sub>2</sub> complex at pH 7.4 and the length of the zinc hook peptide. Black circles correspond to Hk4-Hk45 with native sequences. The  $K_{12}$  values for Hk14VA, Hk14LA and Hk14VALA mutants were adopted from the previous study.<sup>27</sup> Numbers 1 and 2 refer to two ranges of stability increase as discussed in the text.

**Figure 4.** Backbone structure of the  $\beta$ -hairpin-forming fragment of the zinc hook domain. The colors of the circles correspond to the protection factors obtained for Zn(Hk10)<sub>2</sub> in NMR



deuterium exchange experiments. Blue and red dashed lines represent N-H...O and N-H...S hydrogen bonds, respectively. Details are presented in Tables S4 and S5.

**Figure 5.** The  $pK_a^{SH}$  values of cysteine thiols in metal-free hook peptides determined spectrophotometrically at 25°C and  $I = 0.1$  M.


**Figure 6.** ITC titration of  $Zn^{2+}$  with Hk14 peptide. Upper panel shows calorimetric response (raw injection heats) of 50  $\mu$ M  $Zn^{2+}$  titrated with 1.3 mM Hk14 peptide in 50 mM HEPES, pH 7.4, 25°C (black trace) and blank titration with 1.3 mM Hk14 peptide into buffer (red trace). Lower panel displays the corresponding calorimetric binding isotherm.


**Figure 7.** Entropy-enthalpy relationships of the zinc hook complexes. Entropy-enthalpy compensation (EEC) is observed for peptide series where an increase in peptide length increases the folded structure in the corresponding zinc hook complexes (Hk4-Hk8 and Hk31-Hk45) but not for peptide series that present a similar number of folded structures in the dimeric zinc hook complex, regardless of length (Hk10-Hk27).


**Figure 8.** Thermodynamic and structural contributions to the stability of the zinc hook dimer. The values should be read according to the formula  $\Delta G^\circ = \Delta H^\circ_{Zn-pep} + n_H \Delta H^\circ_{CysH} - T \Delta S^\circ$ .  $Zn(Cys_2)_2$ ,  $\beta$ , and cc refer to the thermodynamic contributions in kcal/mol to the complex formation by the particular structural component of the zinc hook domain:  $Zn^{2+}$  binding motif,  $\beta$ -hairpin and coiled-coil, respectively.
























### Structural element:

E (kcal/mol)

$\text{Zn}(\text{Cys}_2)_2$

$\beta$

cc

$\Delta G^\circ$

-20.4

-5.8

-2.1

$\Delta H^\circ$

6.5

-8.9

-4.5

$\Delta H^\circ_{\text{Zn-pep}}$

-25.1

-4.1

-4.2

$n_{\text{H}}\Delta H^\circ_{\text{CysH}}$

31.6

-4.8

-0.3

$-T\Delta S^\circ$

-26.9

3.1

2.4