

Biodiversity and habitats of polar region polyhydroxyalkanoic acid-producing
bacteria: bioprospection by popular screening methods

Małgorzata Marta Rogala¹, Jan Gawor², Robert Gromadka², Magdalena Kowalczyk³, Jakub
Grzesiak^{1*}

<jgrzesiak@ibb.waw.pl>

¹Department of Antarctic Biology, Institute of Biochemistry and Biophysics, Polish Academy of Sciences
Pawieńskiego 5A, 02-106 Warszawa;

²Laboratory of DNA Sequencing and Oligonucleotide Synthesis, Institute of Biochemistry and Biophysics,
Polish Academy of Sciences, Pawieńskiego 5A, 02-106 Warszawa

³Department of Microbial Biochemistry, Institute of Biochemistry and Biophysics, Polish Academy of Sciences
Pawieńskiego 5A, 02-106 Warszawa

Abstract

Polyhydroxyalkanoates (PHA), the intracellular polymers produced by various microorganisms as carbon and energy storage, are of great technological potential as biodegradable versions of common plastics. PHA-producing microbes are therefore in great demand and a plethora of different environments, especially extreme habitats, have been probed for the presence of PHA-accumulators. However, polar region have been neglected in this regard, probably due to the low accessibility of the sampling material and unusual cultivation regime. Here, we present the results of a screening procedure involving 200 bacterial strains isolated for 25 habitats of both polar regions. Agar-based, microscopy and genetic tests were conducted to elucidate the biodiversity and potential of polar-region PHA-accumulators. Microscopic observation of Nile Red stained cells proved to be the most reliable screening method as it allowed to confirm the characteristic bright orange glow of the Nile Red – PHA complex as well as the typical morphology of the PHA inclusions. Psychrophilic PHA-producers belonged mostly to the *Comamonadaceae* family (Betaproteobacteria) although actinobacterial PHA synthesizers of the families

Microbacteriaceae and *Micrococcaceae* also featured prominently. Glacial and postglacial habitats as well as developed polar region soils were evaluated as promising for PHA-producer bioprospection. This study highlights the importance of psychrophiles as a biodiverse and potent polyhydroxyalkanoate sources for scientific and application-aimed research.

Keywords: psychrophiles, Nile Red, *Comamonadaceae*, glacier, feast/famine regime

Introduction

Polyhydroxyalkanoates (PHA) are a group of intracellular polymers synthesized by a variety of prokaryotic microorganisms. Their primary function is that of a carbon and energy storage to be used in starvation periods. They are produced from the excess carbon present in the cells environment and consist of polyesters of hydroxyalkanoic acids with chains of varying lengths. Key enzymes conducting the polymerization process are the PHA synthases encoded by the *phaC* gene. Currently, there are four classes of this enzyme discovered and described. Amino acid sequence overlap between enzymes representing different classes is moderate at best, albeit their products can be divided into scl-PHA – short side chain length PHAs produced mainly by the actions of the I, III and IV class synthases and mcl-PHAs – medium side chain length PHAs produced mainly by the class II synthases harbored *inter alia* by the *Pseudomonas* genus (Reddy *et al.* 2003; Rehm and Steinbüchel 1999).

Polyhydroxyalkanoic acids are of great technological potential, because they are seen as possible petroleum based plastic substitute: biodegradable, biocompatible and derived from biowaste products (Chee *et al.* 2010). Therefore, PHA producing microbes are highly sought after in terms of quality and quantity of the synthesized polymer (Kumar *et al.* 2020). Screening procedures involve mainly the detection of the PHA granule themselves and/or the detection of genes coding the PHA synthesis pathway enzymes (Kung *et al.* 2007). Staining of the granules with lipophilic dyes is the most popular of the methods, with the fluorescent Nile

Red stain being frequently used to differentiate PHA positive bacterial colonies on a Petri dish or to visualize granules within bacterial cells using an adequately equipped microscope. The PHA positive strains display a bright orange glow when irradiated with UV or green (510-560 nm) light (Spiekermann et al. 1999; Kitamura and Doi 1994). Genetic screening involves mainly the PCR based amplification of a fragment of the *phaC* gene using degenerate primers (Romo et al. 2007).

PHA producing bacteria can be found in a variety of environments, mainly those that experience periodic nutrient limitations, as the granule-stored carbon helps to cope with starvation issues (Alistair et al. 1990). Studies conducted on the survival of PHA-containing bacterial cells reveal the involvement of these polymers in cell recovery from a number of other abiotic stressors (Obruca et al. 2020). Therefore, habitats where the intensity of physical and chemical factors change frequently have the potency to enrich the microbial community in most resilient cells, *inter alia* in PHA producers (Pernicowa et al. 2020). Polar region habitats seem in this regard to be a very promising source of novel PHA synthesizing bacterial strains (Ciesielski et al. 2014; Ayub et al. 2004). Polar and circumpolar regions experience a severe case of seasonal nutrient inputs caused by the occurrence of the yearly polar day/night phenomenon. Furthermore, several other factors (not only low temperatures) such as: water activity shifts, freeze-thaw cycles, intense UV radiation, free radical formation and sudden pH changes also contribute to the harshness of this environment (Varin et al. 2012). Therefore, prospecting for polar region cold-loving bacterial PHA producers might prove very fruitful. Such endeavors have to date been undertaken only in a limited respect, presumably due to the low accessibility of the sampling material and unusual cultivation regime (Koller, 2017).

In this study we present a screening of a 200 bacterial isolates derived from 25 different habitats all located in polar regions of both Arctic and Antarctica done by popular testing methods. The aim of these analyses was to evaluate the factual potential of those

microbes to produce PHAs and additionally to find the most accurate method to do so. Our hypothesis states, that polar region habitats harbor a phylogenetically and metabolically diverse cultivable bacteria communities capable of PHA production at low temperatures.

Materials and methods

Strain cultivation

Bacterial strains used in this study (See Tab. 1 for details) are part of the Central Collection of Strains of Institute of Biochemistry and Biophysics Polish Academy of Sciences. The strains are stored as glycerol stocks at -80°C. They were revived by streaking aliquots of the stocks on an appropriate medium. For the saltwater isolates R3A medium was used containing (g/L): Peptone -1, Tryptone – 1, Yeast extract – 1, Beef extract – 1, Glucose – 1, K_2HPO_4 – 1, NaH_2PO_4 – 0.5, $MgSO_4$ – 0.1, prepared with artificial seawater (g/L): NaCl – 27.5, $MgCl_2 \cdot 6H_2O$ – 5.38, $MgSO_4 \cdot 7H_2O$ – 6.78, KCl – 0.72, $NaHCO_3$ – 0.2, $CaCl_2 \cdot 2H_2O$ – 1.4. For all other strains the R3A medium (agar and broth) was prepared with distilled water. The plates were incubated at 15°C in the dark (Liebher Thermostat Cabinet) for a period of 3 weeks. Strains were checked for purity by repeated transfer to fresh agar plates. Single colonies were picked and transferred to test tubes with 3 mL of cool, sterile R3A broth and incubated for 1 week to attain a saturated bacterial culture. The cultures were the basis for further research.

Agar based screening

The R3A agar plates for PHA screening were prepared according to Spiekermann et al. 1999. Stock solutions of Nile Blue A and Nile Red dyes were prepared in DMSO and added to the agar *post* autoclaving to give a final concentration of 0.5 µg dye per mL of medium. Strains were drop plated (5 µL) from the active culture onto the plates, with controls on R3A agar without the dyes. After 1 week of incubation at 15°C the plates were exposed to

ultraviolet light (312 nm) and the drop-spots were examined for the bright orange fluorescence indicative for PHA presence. Those exhibiting the glow were scored as positive.

DNA-based analysis

For DNA extraction from bacterial cells 100µL of the liquid culture was centrifuged at 12000 rpm for 3min. in a 2 mL Eppendorf type tube and the pellet was suspended in 200µL of sterile MiliQ water and a small amount (approx. 10% w/v) of Chelex100 resin and sharp garnet sand were added. The suspension was further amended with 1.5µL of lysozyme solution (10 mg/mL) and incubated at 37°C for 2.5h. Then 10µL of a 10% SDS solution was added to the suspension and the tubes were placed in a Qiagen Retsch TissueLyser II for 5 min. at 39Hz – 1800 oscillation per min. The tubes were centrifuged briefly, amended with 1µL proteinase K solution and incubated at 55°C for 1h. After centrifugation at 12000 rpm for 3min. the DNA in the supernatant was purified using the Clean-up Concentrator kit (A&A Biotechnology) according to manufacturer's protocol. Yield and purity of the extracted DNA was checked in a NanoPhotometer® NP80 (Implen).

Amplification of 16S rRNA gene fragment was performed using universal primers 27F and 1492R (Lane 1991). PCR amplification reaction conditions were as follows: 1 min of 95 °C initial denaturation followed by 30 cycles of 95 °C for 15 s, 55 °C annealing for 15 s and elongation 72 °C for 1min and 30 seconds, using DreamTaq polymerase (Thermo Scientific-Fermentas). Obtained PCR products (~1500 bp for 16S rRNA gene fragment) were checked on 0,8% agarose gel and purified using Clean-up Concentrator kit (A&A Biotechnology). 16S rRNA gene amplicons were sequenced using the 27F 16S rRNA gene primer with the use of BigDye Terminator v.3.1 chemistry and ABI3730xl genetic analyzer at the DNA Sequencing Laboratory (Institute of Biochemistry and Biophysics PAS).

Amplification of *phaC* gene fragment was performed using primers G-D (5'GTGCCGCCSYRSATCAACAAGT3') and G-1R

(5'GTTCCAGWACAGSAKRTCAGAA3') by Romo et al. 2007 targeting both the type I and type II synthase gene. PCR amplification reaction conditions were as follows: 1 cycle of 94°C for 10 min, 60°C for 2 min and 72°C for 2 min followed by 40 cycles at 94°C for 20, 55.5°C for 45 sec and 72°C for 1 min and a final cycle at 72°C for 5 min using DreamTaq polymerase (Thermo Scientific-Fermentas). Obtained PCR products (~550 bp for *phaC* gene fragment) were checked on 0,8% agarose gel. Samples containing the desired length product were scored as positive.

Feast-famine regime implementation

Three types of PHA-inducing media (PIM) were introduced to create a feast/famine regime based on lowering the nitrogen to carbon ratio. The basis was a 7-fold diluted (0.5g/L) R3A medium amended with several carbon sources. PIM1 contained carbon sources that were unrelated to PHA structure: glucose (2.5g/L), sodium lactate (2.5g/L), glycerol (2.5g/L) and sodium acetate (1g/L). PIM2 contained short-chain fatty acids: sodium butyrate and sodium valerate (3.5g/L each). PIM3 contained a mix of medium-chain fatty acids derived from the ultrasonic-initiated saponification of coconut oil (Mercantili et al. 2014). 10g of coconut oil (generic brand) was amended with 30 mL of 1.5M NaOH (in excess) in a 50 mL Falcon-type tube, heated in a water bath at 50°C until the oil completely melted. The ingredients were then shaken for 5 min on a Tornado Vortexer until a homogenic emulsion was achieved. Then the tube was placed in a VWR Ultrasonic Cleaner USC-TH set to 50°C and sonicated at 40kHz for 45 min to facilitate the saponification reaction. The tube was placed in a thermostat at 37°C for 6 weeks to allow the saponification reaction to proceed to completion. The saponified oil was dissolved in ddH₂O, pH was adjusted to 5.0 with HCl upon which the free fatty acids precipitated. The suspension was sedimented by centrifugation (10 min, 8000 rpms) and washed three times with ddH₂O to remove the excess alkali and the glycerol byproduct. Free fatty acids were air-dried and suspended in water at a concentration of 7g/L with Tween80

(0.1g/L). The mixture was emulsified by heating to 45°C and vigorous shaking. All media were adjusted to pH 7.2.

Bacterial cultures on R3A broth were harvested by centrifugation (0.5 mL) in separate Eppendorf tubes, resuspended in 3ml of each of the PIMs and incubated at 14°C for 1 week. Resulting suspension were subjected to microscopy based screening.

Nile red staining and microscopy

Bacterial suspensions were washed with a washing buffer (NaCl -9 g/L, methanol 10 g/L, Tween80 – 50 µg/L, tertrasodium pyrophosphate – 2.6 g/L) to remove media components and slightly perforate cell walls for dye penetration. Bacterial cells were resuspended in 0.9% saline and stained with Nile Red in DMSO (80µg/mL) to give a final concentration of 3.1 µg/mL for 30 min (Zuriani et al. 2013). Stained bacterial suspensions were trapped under a microscope slide and observed under 1000x magnification with green (510-560 nm) and blue light (450-490nm) excitation on a Nikon E-200 microscope with a 100 W Hg lamp and 100× CFI 60 oil immersion objective, with a digital DS-Fi3 high-definition color microscope camera equipped with a 5.9 megapixel CMOS image sensor and a filter block of wavelengths: EX 330–380, DM 400, BA 420.

Data analysis

Calculations charts and graphs were made in Excel (MS Office for Windows). Simple

Matching Coefficient was calculated using the following formula: $SMC = \frac{M_{00} + M_{11}}{M_{00} + M_{01} + M_{10} + M_{11}}$

where: M_{00} - total number of attributes where both have a value of 0, M_{11} - total number of attributes where both have a value of 1, M_{01} and M_{10} - total number of attributes where one has a value of 1 and the other a value of 0. 16S rRNA gene fragments were identified using the Blastn algorithm with the ‘sequences from type material’ option, identifying the strain as the closest match to validly described species. Phylogenetic trees were made using the Mega-X software. Sequences were deposited under the accession numbers: MT585825-MT586024.

Results

Polar region strains screened with the Nile Red-amended agar technique displayed varying shades of the expected bright orange fluorescence. Some strains displayed a fluorescence with a pinkish hue (like *Polaromonas* sp. 1701), some with a more yellowish hue (like *Chrysoebacterium* sp. 966). In others, like the *Rhodanobacter* sp. 2793 only a faint trace of the orange fluorescence could be observed at the rim of the drop-growth (Fig. 1A). Several strains displayed growth retardation on the Nile Red amended agar in comparison to the Nile Red free control. Agar with the Nile Blue A dye did not yield any specific fluorescence. Amplification of the *phaC* gene resulted in the single, specific 550bp long fragment like in the case of *Polaromonas* sp. 1701 or in multiple fragments, including the specific one (Fig. 1B). Microscopic observations of the Nile Red – stained cells observed under green light excitation in most cases confirmed the orange-fluorescing intracellular granules with a clearly defined morphology thereof. The outline of cells without PHA granules was also visible. Most strains belonging to the Bacteroidetes phylum, like *Flavobacterium* sp. 1052 and also marine bacteria (*inter alia* *Psychromonas* sp. 1212) displayed an orange fluorescence of the whole cell, without any granule-characteristic morphology.

Examined isolated belonged to four phyla: Proteobacteria (fig. 2), Actinobacteria, Firmicutes and Bacteroidetes (fig. 3). Proteobacterial isolates belonged to three classes: Alphaproteobacteria (n=8) consisting mostly of *Sphingomonas* sp. strains. Betaproteobacteria were more numerous consisting of 45 isolates clustered within roughly three families, the most important being the *Comamonadaceae* (n=27), with *Polaromonas* as the most frequent genus. Gammaproteobacterial isolates were the most numerous (n=53) and most diverse (>7 families). The family with the most members were the *Pseudomonadaceae* (n=27) consisting exclusively of the *Pseudomonas* genus (fig. 2). Actinobacteria (n=43), Firmicutes (n=6) and

Bacteroidetes (n=45) were less frequently isolated than Proteobacteria. Actinobacterial strains belonged mostly to two families: *Microbacteriaceae* (n=22) and *Micrococcaceae* (n=17). *Microbacteriaceae* consisted mostly of three genera: *Salinibacterium*, *Cryobacterium* and *Glacihabitans*. The *Micrococcaceae* consisted exclusively of two genera: *Arthrobacter* and *Paeniglutamicibacter*. The Bacteroidetes isolates were mostly members of the *Flavobacteriaceae* family (N=36).

To examine the reproducibility of the results obtained by the different methods a Simple Matching Coefficient was calculated between each of the binary data sets, taking both negative and positive responses into account (Fig. 2). As the microscopy-based method was evaluated as the least bias-prone, strains were included into the PHA-positive group (POS) when granules were present after cultivation on either of the liquid media used (R3A, PIM1,2,3). None of the methods used showed a total result overlap. For the ‘All strains’ group the highest overlap was achieved for the *phaC* gene detection and microscopy method for R3A medium cultured strains (CON - constitutive PHA producers) – 0.67. Low overlap values were achieved for the agar plate screening method (0.45-0.59). For Gram-negative bacteria this was also apparent with high overlap of the *phaC* detection with strains scored as positive (0.66). Gram-positive bacteria at the other hand had low overlap (0.34) of the *phaC* detection scores with positive strains but high with microscopy based method for constitutive producers. The plate method displayed in the case of Gram-positive bacteria greater overlap with other methods (0.65 – 0.67). Proteobacteria displayed low to moderate overlap values, especially low for the agar plate method (0.35 - 0.50). Highest overlap value was achieved for the *phaC* detection and PHA positives. Actinobacteria displayed relatively high overlap between *phaC* detection and microscopy method for constitutive producers (0.78) and low when the overlap was calculated for PHA positive strains. Bacteroidetes displayed moderate

to high overlap of tested methods with highest between *phaC* detection and microscopy granule detection after cultivation on R3A broth (Fig. 2.).

Percentage of PHA positive strains were based on microscopy observations (Fig. 5). For all examined strains a value of 63% of PHA-granule positive were achieved. Within the phylum-rank group high values were scored for Proteobacteria (75%) and Actinobacteria. Bacteroidetes scored only 27% PHA-positive strains. Betaproteobacteria and Gammaproteobacteria scored 91% and 60% positive strains respectively. Within the family level groups (where $n > 10$) the *Comamonadaceae* displayed highest values (96%) whereas the *Flavobacteriaceae* the lowest (28%). The *Alcaligenaceae*, *Pseudomonadaceae*, *Microbacteriaceae* and *Micrococcaceae* remained within a 70-80% threshold (Fig. 5). Half of the PHA-producers (within all examined strains) were constitutive granule producers, whereas the other half had to be induced with a high C/N ratio. Highest constitutive producer percentage were found belonging to the *Alcaligenaceae* and *Comamonadaceae* families within the Betaproteobacteria class (86 and 73% respectively), whereas the *Microbacteriaceae* (Actinobacteria) and *Pseudomonadaceae* (Gammaproteobacteria) displayed high percentage of strains that produced PHA only upon induction. The most successful induction medium was the PIM1, inducing PHA-formation on 71% of all of the non-constitutive PHA producers. It was the highest scoring induction medium for all groups except for the Betaproteobacteria and *Alcaligenaceae*, where PIM2 was more successful in inducing PHA formation. PIM3 medium containing medium-length fatty acid salts was the most successful in PHA induction of the *Pseudomonadaceae* family members.

The inducible PHA-granule accumulating strains usually produced the storage material on more than one type of induction media (Fig. 6). PIM1 and PIM2 shared the greatest number of induced strains (23) whereas PIM2 and PIM3 the lowest (13). Twelve strains managed to produce PHAs on all three induction media.

Strains were grouped according to the type of environment they were isolated from (Fig. 7). Glacial environments harbored the most PHA-positive strains (91%), 52% of which were constitutive granule accumulators. High contribution values of PHA positive strains were also apparent for soil-associated (pedogenic) habitats (78%) but also post-glacial deposits (67%). Marine and animal-influenced habitats contained mostly non-PHA producing strains, albeit the PHA producing minority had to be induced to accumulate PHA (>70%). Almost half of the freshwater habitat isolated strains had PHA synthesis abilities, 67% of which did it without the high carbon to nitrogen ratio induction.

Discussion

Evaluating screening methods

Popular screening methods (both physiology- and gene based) displayed a varying degree of accuracy when dealing with polar region-derived bacterial strains. The plate method yielded results that were hard to evaluate. Most difficult to interpret were pigmented strains, which was noted also by other researchers (Higuchi-Takeuchi et al. 2016). In the case of *Rhodanobacter* sp. isolates the presence of the pigment seemed to extinguish the Nile Red-stained granule fluorescence. *Rhodanobacter* isolates often produce xanthomonadin-like pigments, which absorb wavelengths in the blue-green range, those that excite the granule-bound Nile Red dye (He et al. 2020; Rajagopal et al. 1997; Zuriani et al. 2013). Drop-growth of most Bacteroidetes members (mostly *Flavobacterium* spp. and *Chryseobacterium* spp. isolates) and several Proteobacteria of marine origin (e.g. *Psychromonas* sp.) displayed strong orange fluorescence on the Nile Red amended agar. This was checked by microscopy analysis, where whole cells had an orange glow in green light excitation. Flavobacteria of polar origin produce substantial amounts of branched and unsaturated fatty acids to maintain membrane fluidity (Králová 2017), whereas marine psychrophiles produce polyunsaturated fatty acid for the same purpose (Russel and Nichols,

1999). Alonso and Mayzard (1999) state, that those fatty acids belong to the polar lipid group and their fluorescence after Nile Red staining is in the range of 610 nm (orange). A feature of the plate based-screening, besides color evaluation difficulty, was also visible growth retardation on the Nile Red amended agar in comparison to the Nile Red-free control. Presumably, strains that originate from polar regions, where there is low anthropogenic impact and low nutrient concentrations, are more sensitive to xenobiotics, like the diluent used (DMSO) or the dye itself (Znój et al. 2017; Speakermann et al. 1999). Surprisingly, agar amended with the Nile Blue A dye did not yield any specific fluorescence. Presumably the low temperature or the pH of the medium hampered its oxidation to the active derivative (Nile Red) (Greenspan et al. 1985). Genetic screening methods have an advantage over the metabolism based ones as they do not rely on granule presence, scoring however the strains that are being examined only as potential producers. The primer set designed by Romo et al. (2011) targets type I and II synthase genes. Amplification of the *phaC* fragment using those primers frequently yielded multiple bands with or without the 550 bp specific band during the screening of polar region bacteria. This could be explained either by non-specific binding of the degenerate primers or the existence of other copies of the gene in close proximity to each other, simultaneous carrying of two types of synthases or other unusual PHA synthase related genes. Recent studies (Tan et al. 2020) have shown, that *Pseudomonas* spp. isolated from Antarctica carried two classes of synthases (I and II) and some *Janthinobacterium* sp. isolates contained genes of an usual PHA synthase (provisionally named class V) besides the class I synthase. Sequencing of the genomes of several of strains used in this study (data not shown) indicates presence of a variety of genes identified as polyhydroxyalkanoic acid synthases seeded throughout the genome. Furthermore, genome sequencing revealed the presence of type III and IV synthase genes in some of the examined strains. Those genes were not detected by the primers used and currently there is no wide-range primer sets to detect them

(Montenegro et al. 2017). Microscopy observation of the Nile Red stained cells also required cautious examination. However, microscopy-based approach has proven the most accurate and the most reliable down the line, as it detected active PHA accumulators and granule presence was not only confirmed by the characteristic fluorescence but also by their morphology (Wältermann and Steinbüchel 2005). As already mentioned marine strains and those belonging to the Bacteroidetes phylum displayed a whole-cell red fluorescence in green light excitation. Those were scored as negative for PHA accumulation as the granules in native producers tend not to exceed 40% of the cells volume (Mravec et al. 2016). Further advantage of the microscopy-based approach was that the cells were stained *post* growth and PHA production so there were no issues with proliferation-hampering effect of the dye or the carrier diluent (DMSO). Furthermore, the feast-famine regime could easily be incorporated into the method to further increase the number of PHA-positive strains.

Comparison of the data with the Simple Matching Coefficient clearly showed that each method produced different results for the same sample group. Assuming that the microscopic method is the most accurate one and considering the results of the feast/famine regime, several observations can be made. The plate-based method displayed low overlap with the microscopy method in Proteobacteria, suggesting that in this group the agar plate approach potentially introduces a heavy bias. In Actinobacteria *phaC* detection method showed high overlap with the microscopy-based technique only for constitutive producers and low when also inducible producers were taken into account. This points towards failed amplification of most of the actinobacterial *phaC* gene fragment. The reason being, that the primers were designed mostly on the basis of proteobacterial sequences or that polar region actinobacterial genes represent a novel quality among PHA synthases.

Biodiversity of psychrophilic PHA producers

324 The most frequently isolated polar region PHA producing strains belonged mostly to
325 the *Comamonadaceae* family. Members of this family have been frequently recognized as a
326 vital part of PHB – producing consortia enriched from a mixed microbial culture under
327 different feeding conditions (Dai et al. 2015; Cavaillé et al. 2016). Furthermore, members of
328 the *Comamonadaceae* like *Rhodoferax* spp. and *Polaromonas* spp. have been found to
329 produce PHA in sea ice and cold waters of the northern Baltic Sea (Pärnänen et al. 2015).
330 *Polaromonas* spp. are a frequent resident of glacial Arctic and Antarctic habitats and are
331 suspected to be involved in symbiosis with eukaryotic algae (Gawor et al. 2016). This is in
332 line with the conclusions drawn by Kadouri about the pivotal role of PHAs in bacterial
333 symbionts of eukaryotic organisms. Especially surprising is the constitutive feature of the
334 betaproteobacterial PHA synthesis. The need for constant granule presence might be dictated
335 by the sudden changes in nutrient availability or intensity of metabolism-challenging factors
336 as the PHAs are thought to protect extremophiles from environmental stressors (Koller 2017).
337 Gammaproteobacteria, especially the *Pseudomonadaceae* family members produced PHA
338 mostly under nitrogen limiting conditions. Those bacteria are a known group of
339 opportunitrophs, displaying a large variety of adaptive features like fast growth and wide
340 range of substrate utilization and lytic enzyme production (Prieto et al. 2016), so PHA
341 production may be an asset only in extremely harsh conditions (Ayub et al. 2009).
342 Actinobacteria were second in terms of isolate numbers of active PHA producers revealed in
343 this study. A vast majority of actinobacterial strains produced PHA only upon induction with
344 high C/N ratio, most notably the members of the *Microbacteriaceae* family (*Cryobacterium*
345 for eg.). *Micrococcaceae* were more evenly divided among constitutive and inducible PHA
346 producers (eg. *Arthrobacter* spp.). Information on members of those families as PHA
347 producers is limited, however polar region Actinobacteria were described as occupying
348 diverse niches (Zdanowski et al. 2013), so the scenario presented earlier for

Pseudomonadaceae might also be applicable. As reported by Matias et al. 2009 Actinobacteria tend to produce PHAs of unusual chemistry, therefore actinobacterial isolates derived from polar region materials might represent an untapped source of novel, biotechnologically relevant polyhydroxyalkanoates.

PHA producers and their habitat

Polar region habitat types differed quite substantially in cultivable bacterial PHA accumulators' content. The zoogenic habitat type which was represented by sea birds' (penguins and little auks) nesting sites and marine habitats represented mainly by decaying seaweed heaps displayed the lowest contribution of PHA producers. Seemingly, polar nutrient-rich substratum (Grzesiak et al. 2020, Zdanowski et al. 2005) does not promote species that diverge energy and resources towards carbon-storage material synthesis (Pratt et al. 2012; Wang and Yu, 2000). Furthermore, even among isolates displaying this ability, its expression had to be triggered by nitrogen starvation with simultaneous carbon surplus. Therefore, constant high nutrient availability negatively influences the contribution of PHA producers to the community also in polar regions. Contrasting in this respect were the highly oligotrophic glacial habitats. Vital nutrients, especially nitrogen and phosphorus are scarce in those sites (Grzesiak et al. 2015a; Grzesiak et al. 2015b). Over 90% contribution of PHA positive isolates among glacier-derived bacteria was therefore not surprising. Presumably, labile carbon surplus is present only during the short summer season (Telling et al. 2014), therefore PHA production is needed in regards of carbon deficit survival. Among glacier-derived PHA⁺ strains there was an even distribution of constitutive and inducible PHA producers, indicating different mechanisms for coping with extreme glacial conditions. As mentioned before, some bacterial groups might use the granules as an abiotic stress-resistance feature, whereas others might rely on different mechanisms, not related to PHA metabolism like pigment synthesis (Dieser et al. 2010). Soils of varying complexity (developed and

postglacial) shared similar characteristics to each other in terms of percentage of PHA-producing strains and their ability to constitutive or inducible granule accumulation. Heterogeneity of such habitat enables an establishment of a multitude of different niches, even among closely related microorganisms (Gittel et al. 2014), therefore the fairly even distribution between non-producers, constitutive producers and inducible producers. Polar region freshwater habitats are strongly affected by seasonal changes in nutrient quantity and quality, creating a natural feast/famine regime (Crump et al. 2003). Consequently, high altitude Himalayan lakes were recently proclaimed as “bioplastics reservoir” by Kumar and associates (2018).

Conclusions

Polar region bacteria present a novel and potent source of PHA-producing microorganisms. Microscopic observation of Nile Red stained cells amended with a feast/famine regime implementation using different carbon sources was proven as the most reliable screening method. Caution is advised when evaluating members of the Bacteroidetes phylum and polar region marine bacteria as they often give false positive results in the Nile Red dye involving approach. Members of the *Comamonadaceae* family were the most numerous PHA producers by percentage among the 200 examined strains. Most abundant in cultivable PHA accumulators were glacier associated habitats of both polar regions, followed by well-developed soils and postglacial deposits. Nonetheless, to further expand the topic of polar region PHA producers additional investigations are in order: a culture independent approach of the native source material and an in depth laboratory analysis of bacterial strains.

Acknowledgments

This work was supported by the National Science Center, Poland (Grant 2017/24/C/NZ9/00232). Samples and data were obtained thanks to the scientific facilities of the Polish Antarctic Station ARCTOWSKI and the Polish Polar Station HORNSUND.

399 **References**

- 400 Anderson, A. J., & Dawes, E. A. (1990). Occurrence, metabolism, metabolic role, and
401 industrial uses of bacterial polyhydroxyalkanoates. *Microbiology and Molecular Biology*
402 *Reviews*, 54(4), 450-472.
- 403 Alonzo, F., & Mayzaud, P. (1999). Spectrofluorometric quantification of neutral and polar
404 lipids in zooplankton using Nile red. *Marine chemistry*, 67(3-4), 289-301.
- 405 Ayub, N. D., Pettinari, M. J., Ruiz, J. A., & López, N. I. (2004). A polyhydroxybutyrate-
406 producing *Pseudomonas* sp. isolated from Antarctic environments with high stress
407 resistance. *Current microbiology*, 49(3), 170-174.
- 408 Ayub, N. D., Tribelli, P. M., & López, N. I. (2009). Polyhydroxyalkanoates are essential for
409 maintenance of redox state in the Antarctic bacterium *Pseudomonas* sp. 14-3 during low
410 temperature adaptation. *Extremophiles*, 13(1), 59-66.
- 411 Cavaillé, L., Albuquerque, M., Grousseau, E., Lepeuple, A. S., Uribe Larrea, J. L., Hernandez-
412 Raquet, G., & Paul, E. (2016). Understanding of polyhydroxybutyrate production under
413 carbon and phosphorus-limited growth conditions in non-axenic continuous
414 culture. *Bioresource technology*, 201, 65-73.
- 415 Chee, J. Y., Yoga, S. S., Lau, N. S., Ling, S. C., Abed, R. M., & Sudesh, K. (2010).
416 Bacterially produced polyhydroxyalkanoate (PHA): converting renewable resources into
417 bioplastics. *Current research, technology and education topics in Applied Microbiology and*
418 *Microbial Biotechnology*, 2, 1395-1404.
- 419 Ciesielski, S., Górniak, D., Możejko, J., Świątecki, A., Grzesiak, J., & Zdanowski, M. (2014).
420 The diversity of bacteria isolated from Antarctic freshwater reservoirs possessing the ability
421 to produce polyhydroxyalkanoates. *Current microbiology*, 69(5), 594-603.
- 422 Crump, B. C., Kling, G. W., Bahr, M., & Hobbie, J. E. (2003). Bacterioplankton community
423 shifts in an arctic lake correlate with seasonal changes in organic matter source. *Appl.*
424 *Environ. Microbiol.*, 69(4), 2253-2268.
- 425 Dai, J., Gliniewicz, K., Settles, M. L., Coats, E. R., & McDonald, A. G. (2015). Influence of
426 organic loading rate and solid retention time on polyhydroxybutyrate production from hybrid
427 poplar hydrolysates using mixed microbial cultures. *Bioresource technology*, 175, 23-33.
- 428 Diesler, M., Greenwood, M., & Foreman, C. M. (2010). Carotenoid pigmentation in Antarctic
429 heterotrophic bacteria as a strategy to withstand environmental stresses. *Arctic, Antarctic, and*
430 *Alpine Research*, 42(4), 396-405.
- 431 Gawor, J., Grzesiak, J., Sasin-Kurowska, J., Borsuk, P., Gromadka, R., Górniak, D., ... &
432 Zdanowski, M. K. (2016). Evidence of adaptation, niche separation and microevolution
433 within the genus *Polaromonas* on Arctic and Antarctic glacial surfaces. *Extremophiles*, 20(4),
434 403-413.
- 435 Gittel, A., Bárta, J., Kohoutová, I., Mikutta, R., Owens, S., Gilbert, J., ... & Lashchinskiy, N.
436 (2014). Distinct microbial communities associated with buried soils in the Siberian
437 tundra. *The ISME journal*, 8(4), 841-853.

438 Greenspan, P., Mayer, E. P., & Fowler, S. D. (1985). Nile red: a selective fluorescent stain for
439 intracellular lipid droplets. *The Journal of cell biology*, 100(3), 965-973.

440 Grzesiak, J., Górniak, D., Świątecki, A., Aleksandrak-Piekarczyk, T., Szatraj, K., &
441 Zdanowski, M. K. (2015). Microbial community development on the surface of Hans and
442 Werenskiold Glaciers (Svalbard, Arctic): a comparison. *Extremophiles*, 19(5), 885-897.

443 Grzesiak, J., Zdanowski, M. K., Górniak, D., Świątecki, A., Aleksandrak-Piekarczyk, T.,
444 Szatraj, K., ... & Nieckarz, M. (2015). Microbial community changes along the ecology
445 glacier ablation zone (King George Island, Antarctica). *Polar Biology*, 38(12), 2069-2083.

446 Grzesiak, J., Kaczyńska, A., Gawor, J., Żuchniewicz, K., Aleksandrak-Piekarczyk, T.,
447 Gromadka, R., & Zdanowski, M. K. (2020). A smelly business: Microbiology of Adélie
448 penguin guano (Point Thomas rookery, Antarctica). *Science of The Total Environment*,
449 136714.

450 He, Y. W., Cao, X. Q., & Poplawsky, A. R. (2020). Chemical Structure, Biological Roles,
451 Biosynthesis and Regulation of the Yellow Xanthomonadin Pigments in the Phytopathogenic
452 Genus Xanthomonas. *Molecular Plant-Microbe Interactions*, 33(5), 705-714.

453 Higuchi-Takeuchi, M., Morisaki, K., & Numata, K. (2016). A screening method for the
454 isolation of polyhydroxyalkanoate-producing purple non-sulfur photosynthetic bacteria from
455 natural seawater. *Frontiers in microbiology*, 7, 1509.

456 Kadouri, D., Jurkevitch, E., Okon, Y., & Castro-Sowinski, S. (2005). Ecological and
457 agricultural significance of bacterial polyhydroxyalkanoates. *Critical reviews in*
458 *microbiology*, 31(2), 55-67.

459 Kitamura, S., & Doi, Y. (1994). Staining method of poly (3-hydroxyalkanoic acids) producing
460 bacteria by Nile blue. *Biotechnology techniques*, 8(5), 345-350.

461 Koller, M. (2017). Production of polyhydroxyalkanoate (PHA) biopolyesters by
462 extremophiles. *MOJ Polymer Science*, 1(2), 1-19.

463 Králová, S. (2017). Role of fatty acids in cold adaptation of Antarctic psychrophilic
464 Flavobacterium spp. *Systematic and applied microbiology*, 40(6), 329-333.

465 Kumar, V., Thakur, V., Kumar, S., & Singh, D. (2018). Bioplastic reservoir of diverse
466 bacterial communities revealed along altitude gradient of Pangi-Chamba trans-Himalayan
467 region. *FEMS microbiology letters*, 365(14), fny144.

468 Kumar, V., Kumar, S., & Singh, D. (2020). Microbial polyhydroxyalkanoates from extreme
469 niches: Bioprospection status, opportunities and challenges. *International Journal of*
470 *Biological Macromolecules*, 147, 1255-1267.

471 Kung, S. S., Chuang, Y. C., Chen, C. H., & Chien, C. C. (2007). Isolation of
472 polyhydroxyalkanoates-producing bacteria using a combination of phenotypic and genotypic
473 approach. *Letters in applied microbiology*, 44(4), 364-371.

474 Lane, D. J. (1991). 16S/23S rRNA sequencing In: Stackebrandt E, Goodfellow M, editors.
475 Nucleic acid techniques in bacterial systematics.

476 Matias, F., Bonatto, D., Padilla, G., Rodrigues, M. F. D. A., & Henriques, J. A. P. (2009).
477 Polyhydroxyalkanoates production by actinobacteria isolated from soil. *Canadian journal of*
478 *Microbiology*, 55(7), 790-800.

479 Mercantili, L., Davis, F., Collyer, S. D., & Higson, S. P. (2015). Ultrasonic modification of
480 the viscosity of vegetable, mineral and synthetic oils—effects of nucleating agents and
481 free-radical scavengers. *Lubrication Science*, 27(3), 177-191.

482 Montenegro, E. M. D. S., Delabary, G. S., Silva, M. A. C. D., Andreote, F. D., & Lima, A. O.
483 D. S. (2017). Molecular diagnostic for prospecting polyhydroxyalkanoate-producing
484 bacteria. *Bioengineering*, 4(2), 52.

485 Mravec, F., Obruca, S., Krzyzanek, V., Sedlacek, P., Hrubanova, K., Samek, O., ... &
486 Nebesarova, J. (2016). Accumulation of PHA granules in *Cupriavidus necator* as seen by
487 confocal fluorescence microscopy. *FEMS microbiology letters*, 363(10).

488 Obruca, S., Sedlacek, P., Koller, M., Kucera, D., & Pernicova, I. (2018). Involvement of
489 polyhydroxyalkanoates in stress resistance of microbial cells: Biotechnological consequences
490 and applications. *Biotechnology advances*, 36(3), 856-870.

491 Pärnänen, K., Karkman, A., Virta, M., Eronen-Rasimus, E., & Kaartokallio, H. (2015).
492 Discovery of bacterial polyhydroxyalkanoate synthase (PhaC)-encoding genes from seasonal
493 Baltic Sea ice and cold estuarine waters. *Extremophiles*, 19(1), 197-206.

494 Pernicova, I., Novackova, I., Sedlacek, P., Kourilova, X., Koller, M., & Obruca, S. (2020).
495 Application of osmotic challenge for enrichment of microbial consortia in
496 polyhydroxyalkanoates producing thermophilic and thermotolerant bacteria and their
497 subsequent isolation. *International Journal of Biological Macromolecules*, 144, 698-704.

498 Pratt, S., Werker, A., Morgan-Sagastume, F., & Lant, P. (2012). Microaerophilic conditions
499 support elevated mixed culture polyhydroxyalkanoate (PHA) yields, but result in decreased
500 PHA production rates. *Water Science and Technology*, 65(2), 243-246.

501 Prieto, A., Escapa, I. F., Martínez, V., Dinjaski, N., Herencias, C., de la Peña, F., ... &
502 Revelles, O. (2016). A holistic view of polyhydroxyalkanoate metabolism in *Pseudomonas*
503 *putida*. *Environmental microbiology*, 18(2), 341-357.

504 Rajagopal, L., Sundari, C. S., Balasubramanian, D., & Sonti, R. V. (1997). The bacterial
505 pigment xanthomonadin offers protection against photodamage. *FEBS letters*, 415(2), 125-
506 128.

507 Reddy, C. S. K., Ghai, R., & Kalia, V. (2003). Polyhydroxyalkanoates: an
508 overview. *Bioresource technology*, 87(2), 137-146.

509 Rehm, B. H., & Steinbüchel, A. (1999). Biochemical and genetic analysis of PHA synthases
510 and other proteins required for PHA synthesis. *International journal of biological*
511 *macromolecules*, 25(1-3), 3-19.

512 Revelo Romo, D. M., Grosso, M. V., Moreno Solano, N. C., & Montoya Castaño, D. (2007).
513 A most effective method for selecting a broad range of short and medium-chain-length
514 polyhydroxyalkanoate producing microorganisms. *Electronic Journal of Biotechnology*, 10(3),
515 348-357.

- Russell, N. J., & Nichols, D. S. (1999). Polyunsaturated fatty acids in marine bacteria—a dogma rewritten. *Microbiology*, 145(4), 767-779.
- Spiekermann, P., Rehm, B. H., Kalscheuer, R., Baumeister, D., & Steinbüchel, A. (1999). A sensitive, viable-colony staining method using Nile red for direct screening of bacteria that accumulate polyhydroxyalkanoic acids and other lipid storage compounds. *Archives of microbiology*, 171(2), 73-80.
- Tan, I. K. P., Foong, C. P., Tan, H. T., Lim, H., Zain, N. A. A., Tan, Y. C., ... & Sudesh, K. (2020). Polyhydroxyalkanoate (PHA) synthase genes and PHA-associated gene clusters in *Pseudomonas* spp. and *Janthinobacterium* spp. isolated from Antarctica. *Journal of biotechnology*.
- Telling, J., Anesio, A. M., Tranter, M., Fountain, A. G., Nylen, T., Hawkings, J., ... & Wadham, J. L. (2014). Spring thaw ionic pulses boost nutrient availability and microbial growth in entombed Antarctic Dry Valley cryoconite holes. *Frontiers in microbiology*, 5, 694.
- Varin, T., Lovejoy, C., Jungblut, A. D., Vincent, W. F., & Corbeil, J. (2012). Metagenomic analysis of stress genes in microbial mat communities from Antarctica and the High Arctic. *Appl. Environ. Microbiol.*, 78(2), 549-559.
- Wältermann, M., & Steinbüchel, A. (2005). Neutral lipid bodies in prokaryotes: recent insights into structure, formation, and relationship to eukaryotic lipid depots. *Journal of bacteriology*, 187(11), 3607-3619.
- Wang, J., & Yu, J. (2000). Kinetic analysis on inhibited growth and poly (3-hydroxybutyrate) formation of *Alcaligenes eutrophus* on acetate under nutrient-rich conditions. *Process Biochemistry*, 36(3), 201-207.
- Zdanowski, M. K., Zmuda, M. J., & Zwolska, I. (2005). Bacterial role in the decomposition of marine-derived material (penguin guano) in the terrestrial maritime Antarctic. *Soil Biology and Biochemistry*, 37(3), 581-595.
- Zdanowski, M. K., Żmuda-Baranowska, M. J., Borsuk, P., Świątecki, A., Górniak, D., Wolicka, D., ... & Grzesiak, J. (2013). Culturable bacteria community development in postglacial soils of Ecology Glacier, King George Island, Antarctica. *Polar Biology*, 36(4), 511-527.
- Znój, A., Chwedorzewska, K. J., Androsiuk, P., Cuba-Diaz, M., Giełwanowska, I., Koc, J., ... & Zmarz, A. (2017). Rapid environmental changes in the Western Antarctic Peninsula region due to climate change and human activity. *Applied Ecology and Environmental Research*, 15(4), 525-539.
- Zuriani, R., Vigneswari, S., Azizan, M. N. M., Majid, M. I. A., & Amirul, A. A. (2013). A high throughput Nile red fluorescence method for rapid quantification of intracellular bacterial polyhydroxyalkanoates. *Biotechnology and bioprocess engineering*, 18(3), 472-478.

Strain numbers	Material of isolation	Place of origin	Environment type	Maintenance medium
2043, 2045, 2047, 2049, 2099, 2102, 2132, 2139, 2157	Little auk (Alle alle) guano	Hornsund Fiord, Spitsbergen Island, Arctic	Zoogenic	R3A agar
966, 967, 968, 969, 970, 972, 973, 974, 975	Adélie penguin guano	Point Thomas Rookery, King George Island, Antarctica	Zoogenic	R3A agar
1198, 1200, 1204, 1206, 1208, 1210, 1212, 1334, 1335, 1336, 1337, 1338, 1339, 1340, 1341, 1343, 2848	Decaying seaweeds	Hornsund Fiord, Spitsbergen Island, Arctic	Marine	R3A agar with artificial sea water
1639, 1640, 1641, 1642, 1643, 1662, 1663, 1664, 1666, 2861	Decaying seaweeds	Admiralty Bay shore, King George Island, Antarctica	Marine	R3A agar with artificial sea water
2529, 2539, 2543, 2556, 2561, 2574, 2579, 2595, 2600	Cryoconite	Hans Glacier, Spitsbergen, Arctic	Glacial	R3A agar
2720, 2722, 2724, 2728	Cryoconite	Werenskiöld Glacier, Spitsbergen, Arctic	Glacial	R3A agar
301, 303, 304, 305, 330, 331, 332, 333, 334	Glacial surface ice	Ecology Glacier, King George Island, Antarctica	Glacial	R3A agar
473, 474, 475, 476, 477, 478, 479, 481, 482	Glacial surface ice	Baranowski Glacier, King George Island, Antarctica	Glacial	R3A agar
803, 805, 806, 807, 808, 809, 810, 811, 812	Plant-free postglacial soil	Ecology Glacier foreland, King George Island, Antarctica	Glacial	R3A agar
989, 990, 991, 992, 993, 994, 995, 996, 997, 998	Postglacial soil with plant debris	Ecology Glacier foreland, King George Island, Antarctica	Glacial	R3A agar
844, 847, 848, 851	Plant-free postglacial soil	Baranowski Glacier foreland, King George Island, Antarctica	Glacial	R3A agar
117, 123, 130, 139, 1258, 1259, 1261	Plant-free postglacial soil	Windy Glacier foreland, King George Island, Antarctica	Glacial	R3A agar
2176, 2180, 2181, 2183, 2186, 2191, 2196, 2199, 2200, 2201, 2203	Arctic tundra soil with moss/lichen debris	Hornsund Fiord, Spitsbergen Island, Arctic	Pedogenic	R3A agar
1044, 1051, 1052, 1053, 1058, 1062, 1065, 1079, 1081	Kettle lake water	Werenskiöld Glacier forefield, Hornsund fiord, Spitsbergen, Arctic	Freshwater	R3A agar
1699, 1700, 1701, 1702, 1703, 1704, 1705, 1706, 1707, 1708, 2031, 2032, 2033, 2034, 2035, 2036, 2042, 2083, 2084, 2954, 2955	Freshwater microbial mat	Jasnorzewski Gardens, King George Island, Antarctica	Freshwater	R3A agar
867, 868, 869, 870, 871, 872,	Air	Point Thomas	Zoogenic	R3A agar

873, 874, 875, 876		Rookery, King George Island, Antarctica		
1160, 1162, 1165, 1166, 1167, 1169, 1170, 1191, 1192	Subglacial water	Subglacial stream, Hans Glacier, Spitsbergen, Arctic	Glacial	R3A agar
P12, 2778, 2779, 2785	<i>Deschampsia antarctica</i> rhizosphere soil	Arctowski Station vicinity, King George Island, Antarctica	Pedogenic	R3A agar
2792, 2793, 2794, 2795, 2797, 2799, 2800, 2860	Ornithogenic soil	Arctowski Station vicinity, King George Island, Antarctica	Pedogenic	R3A agar
2810, 2812, 2813, 2814, 2815, 2816, 2817, 2818, 2819	Moss rhizosphere	Arctowski Station vicinity, King George Island, Antarctica	Pedogenic	R3A agar
2921, 2923, 2924, 2926, 2927, 2928	Supraglacial water	Supraglacial stream, Ecology Glacier, King George Island, Antarctica	Glacial	R3A agar
2743, 2744, 2746, 2747, 2748, 2749, 2750	River water	Ariedalen stream, Spitsbergen, Arctic	Freshwater	R3A agar

Table 1. Information on origin and cultivation of strains used in analysis.




Fig 1. (A) Examples of drop growth fluorescence of bacterial strains in UV light on a R3A agar supplemented with Nile Red (from left to right): *Janthinobacterium* sp. 1167, *Chryseobacterium* sp. 966, *Rhodanobacter* sp. 2793. (B) Examples of PCR-amplification results (bands) of a *phaC* gene fragment using the G-D and G-1R primers by Romo et al. 2007 (left to right): *Polaromonas* sp. 1701 (550bp band), *Acidovorax* sp. 1169 (300 bp and 550 bp band), *Janthinobacterium* sp. 1167 (550 bp and 1000 bp band), (C) Bacterial cell fluorescence in green light excitation (510-560 nm) after Nile Red staining (left to right): *Janthinobacterium* sp. 1167 – “empty” cell, *Janthinobacterium* sp. 1167 – granule-filled cell, *Flavobacterium* sp. 1052 cell, *Psychromonas* sp. 1212 cell.




Fig 2. Phylogenetic tree based on partial 16S rRNA gene sequences belonging to the Proteobacteria phylum lined up with the results from the PHA screening with various methods. The tree was built using the neighbor-joining method. Bootstrap values are indicated at the nodes. Green boxes indicate a positive score. *phaC* – presence of the 550 bp DNA fragment of the *phaC* gene; PLATE – presence of characteristic fluorescence on R3A agar plates with Nile Red after UV exposure; R3A – presence of red-fluorescing granules in Nile Red stained cells cultured on R3A broth; PIM1 - presence of red-fluorescing granules in Nile Red stained cells incubated in PIM1 medium; PIM2 - presence of red-fluorescing granules in Nile Red stained cells incubated in PIM2 medium; PIM3 - presence of red-fluorescing granules in Nile Red stained cells incubated in PIM3 medium.




Fig 3. Phylogenetic tree based on on partial 16S rRNA gene sequences belonging to the Actinobacteria, Firmicutes and Bacteroidetes phylum lined up with the results from the PHA screening with various methods. The tree was built using the neighbor-joining method. Bootstrap values are indicated at the nodes. Green boxes indicate a positive score. *phaC* – presence of the 550 bp DNA fragment of the *phaC* gene; PLATE – presence of characteristic fluorescence on R3A agar plates with Nile Red after UV exposure; R3A – presence of red-fluorescing granules in Nile Red stained cells cultured on R3A broth; PIM1 - presence of red-fluorescing granules in Nile Red stained cells incubated in PIM1 medium; PIM2 - presence of red-fluorescing granules in Nile Red stained cells incubated in PIM2 medium; PIM3 - presence of red-fluorescing granules in Nile Red stained cells incubated in PIM3 medium.

All strains

	PLATE	CON ^M	POS ^M
<i>phaC</i>	0.59	0.67	0.58
	PLATE	0.52	0.45

Proteobacteria

	PLATE	CON ^M	POS ^M
<i>phaC</i>	0.50	0.54	0.65
	PLATE	0.38	0.35

Gram-negative bacteria

	PLATE	CON ^M	POS ^M
<i>phaC</i>	0.55	0.64	0.66
	PLATE	0.47	0.43

Actinobacteria

	PLATE	CON ^M	POS ^M
<i>phaC</i>	0.65	0.79	0.30
	PLATE	0.63	0.51

Gram-positive bacteria

	PLATE	CON ^M	POS ^M
<i>phaC</i>	0.67	0.78	0.34
	PLATE	0.65	0.51

Bacteroidetes

	PLATE	CON ^M	POS ^M
<i>phaC</i>	0.69	0.87	0.69
	PLATE	0.69	0.60

628

629 Fig. 4. Simple Matching Coefficient calculated from the binary (positive/negative) data
630 obtained from each screening method for the following subsets: All strains (n=200), Gram-
631 negative bacteria (n=151), Gram-positive bacteria (n=49), Proteobacteria (n=106),
632 Actinobacteria (n=43), Bacteroidetes (n=45). *phaC* – binary dataset of the presence/absence
633 of the 550 bp DNA fragment of the *phaC* gene; PLATE – binary dataset of the
634 presence/absence of the characteristic fluorescence on R3A agar plates with Nile Red after
635 UV exposure; CON^M – microscopy-obtained (M) binary dataset of the presence/absence of
636 granules after incubation in R3A broth (constitutive producers); POS^M - microscopy-obtained
637 (M) binary dataset of the presence/absence of granules after incubation on ≥ 1 of the liquid
638 media used (R3A, PIM1,2,3).




Fig 5. Percentage of strains (total and within selected taxonomic ranks) expressing the following traits: Positive - presence of red-fluorescing granules (on R3A broth and on PIMs); Negative – absence of red-fluorescing granules in any conditions; Constitutive – presence of red-fluorescing granules in R3A broth; Induced - presence of red-fluorescing granules only after incubation in PIMs; PIM1 - presence of red-fluorescing granules after incubation in PIM1 within the ‘Induced’ group; PIM2 - presence of red-fluorescing granules after incubation in PIM2 within the ‘Induced’ group; PIM3 - presence of red-fluorescing granules after incubation in PIM3 within the ‘Induced’ group.





Fig 6. Venn diagram displaying numbers of strains within the ‘Induced’ group that were PHA positive on different version of PIM. Blue circle – PIM1, green circle – PIM2, orange circle – PIM3.



678

679 Fig 7. Percentage of strains divided according to habitat-type of origin: A – zoogenic habitat,
 680 B – marine habitat, C – glacial habitats, D – pedogenic habitats, E – postglacial habitats, F –
 681 freshwater habitats. Percentage of strains positive for granule presence – green part of
 682 diagram, negative – gray, putative PHA producers within the positives – orange, PIM-
 683 inducible PHA producers within the positives – yellow.