

ANNEXIN1 mediates calcium-dependent systemic defense in Arabidopsis plants upon herbivory and wounding

Jaiana Malabarba^{1,2} , Anja K. Meents^{1,3} , Michael Reichelt⁴ , Sandra S. Scholz³ , Edgar Peiter⁵ , Julia Rachowka⁶ , Dorota Konopka-Postupolska⁶ , Katie A. Wilkins⁷ , Julia M. Davies⁷ , Ralf Oelmüller³  and Axel Mithöfer¹ 

¹Research Group Plant Defense Physiology, Max Planck Institute for Chemical Ecology, Jena 07745, Germany; ²Postgraduate Program in Cell and Molecular Biology, Biotechnology Center, Federal University of Rio Grande do Sul, Porto Alegre, RS 90040-060, Brazil; ³Plant Physiology, Matthias-Schleiden-Institute for Genetics, Bioinformatics and Molecular Botany, Faculty of Biological Sciences, Friedrich-Schiller-University Jena, Jena 07743, Germany; ⁴Department of Biochemistry, Max Planck Institute for Chemical Ecology, Jena 07745, Germany; ⁵Institute of Agricultural and Nutritional Sciences, Faculty of Natural Sciences III, Martin Luther University of Halle-Wittenberg, Halle (Saale) 06120, Germany; ⁶Plant Protein Phosphorylation Laboratory, Institute of Biochemistry and Biophysics, Polish Academy of Sciences, Warsaw 02-106, Poland; ⁷Department of Plant Sciences, University of Cambridge, Cambridge, CB24 6DG, UK

Summary

Author for correspondence:
Axel Mithöfer
Email: amithoef@ice.mpg.de

Received: 12 March 2020
Accepted: 5 February 2021

New Phytologist (2021)
doi: 10.1111/nph.17277

Key words: calcium signaling, herbivory, jasmonates, plant defense, *Spodoptera*, systemic signaling.

- Cellular calcium (Ca) transients are endogenous signals involved in local and systemic signaling and defense activation upon environmental stress, including wounding and herbivory. Still, not all Ca²⁺ channels contributing to the signaling have been identified, nor are their modes of action fully known. Plant annexins are proteins capable of binding to anionic phospholipids and can exhibit Ca channel-like activity. Arabidopsis ANNEXIN1 (ANN1) is suggested to contribute to Ca transport.
- Here, we report that wounding and simulated-herbivory-induced cytosolic free Ca elevation was impaired in systemic leaves in *ann1* loss-of-function plants. We provide evidence for a role of ANN1 in local and systemic defense of plants attacked by herbivorous *Spodoptera littoralis* larvae.
- Bioassays identified ANN1 as a positive defense regulator. *Spodoptera littoralis* feeding on *ann1* gained significantly more weight than larvae feeding on wild-type, whereas those feeding on ANN1-overexpressing lines gained less weight. Herbivory and wounding both induced defense-related responses on treated leaves, such as jasmonate accumulation and defense gene expression. These responses remained local and were strongly reduced in systemic leaves in *ann1* plants.
- Our results indicate that ANN1 plays an important role in activation of systemic rather than local defense in plants attacked by herbivorous insects.

Introduction

Plants are challenged throughout their life by various abiotic and biotic stress factors. These changes in the environment require fast adaptation. Consequently, plants evolved a multi-layered metabolic barrier, composed of mechanical and chemical defenses (Maffei *et al.*, 2012; Mithöfer & Boland, 2012). An attack by herbivorous insects represents a major threat to the plant's survival. In particular, an attack by chewing insects is a combination of plant tissue wounding and application of insect-specific herbivore-associated molecular patterns, mainly present in their oral secretions (OSs; Mithöfer & Boland, 2008; Vadassery *et al.*, 2012b; Kiep *et al.*, 2015). The establishment of chemical defenses to such an insect herbivory is mediated by a network of signaling pathways (including calcium (Ca²⁺) ions, protein phosphorylation, phytohormones,

and reactive oxygen species (ROS) and reactive nitrogen (N) species) that finally initializes synthesis and accumulation of a plethora of defensive metabolites (Seybold *et al.*, 2014; Zebelo *et al.*, 2014). The elevation in cytosolic free Ca, [Ca²⁺]_{cyt}, is one of the earliest signaling events initiated upon the plant's interaction with feeding insects (Maffei *et al.*, 2004; Kiep *et al.*, 2015; Toyota *et al.*, 2018). Jasmonates represent the most important class of wound-induced phytohormones to be activated, with the main components being jasmonic acid (JA) and its biologically active isoleucine conjugate (+)-7-*iso*-jasmonoyl-L-isoleucine (JA-Ile) (Wasternack, 2007; Mithöfer & Boland, 2008, 2012). The connection between jasmonates and [Ca²⁺]_{cyt} is likely mediated by Ca²⁺-sensing proteins. In plants, canonical calmodulins, calmodulin-like proteins (CMLs), calcineurin B-like proteins, and Ca²⁺-dependent protein kinases are good candidates (Swarbreck *et al.*, 2013; Yan

et al., 2018; Mohanta *et al.*, 2019; Tai *et al.*, 2019). In particular, a connection between $[Ca^{2+}]_{cyt}$ and jasmonate signaling has been shown for CML42 and CML37 (DeFalco *et al.*, 2010; Vadassery *et al.*, 2012a,b; Scholz *et al.*, 2014, 2016; Heyer *et al.*, 2018b).

However, upstream of jasmonates, Ca^{2+} -sensing proteins, and $[Ca^{2+}]_{cyt}$ changes, an influx of Ca^{2+} is necessary to cause $[Ca^{2+}]_{cyt}$ elevations. Here, the opening of ion channels localized at the plasma membrane or intracellular compartments is involved. In *Arabidopsis*, the ligand-gated channels glutamate receptor (GLR)-like channels and cyclic nucleotide-gated channels (CNGCs) plus the potentially stretch-activated Ca^{2+} channels reduced hyperosmolality-induced Ca^{2+} increase and MID1-complementing activity are the four main families of plasma membrane Ca^{2+} -permeable channels (Dodd *et al.*, 2010; Yuan *et al.*, 2014). In addition, the vacuolar two-pore channel 1 (TPC1) is localized at the tonoplast (Peiter *et al.*, 2005; Peiter, 2011).

Strikingly, *Arabidopsis* senses local herbivore attack and transmits this information to unwounded vascular-connected systemic leaves through a long-distance signaling system (Mousavi *et al.*, 2013; Kiep *et al.*, 2015). Systemic signaling has also been identified leading to activation of jasmonate accumulation and signaling in distal leaves (Mousavi *et al.*, 2013; Heyer *et al.*, 2018b). Recently, it has been shown that wound-induced electrical signals precede vascular Ca^{2+} fluxes as xylem contact cells and phloem sieve elements function together for leaf-to-leaf electrical signaling (Nguyen *et al.*, 2018). Probably, the systemic electrical signaling is mediated by GLR-type cation channels, because in *glr3.3 glr3.6* double mutants the wound-activated electrical signal propagation, as well as propagation of $[Ca^{2+}]_{cyt}$ signals between leaves, is attenuated (Toyota *et al.*, 2018). In addition, TPC1 was shown to be involved in systemic $[Ca^{2+}]_{cyt}$ elevations (Kiep *et al.*, 2015). This channel trio of GLR3.3, GLR3.6 and TPC1 also operates in local $[Ca^{2+}]_{cyt}$ elevations induced by aphid feeding (Vincent *et al.*, 2017). Very recently, it was demonstrated that a rapidly activated cyclic nucleotide-gated Ca^{2+} channel (CNGC19) also plays a partial role in wounding-induced Ca^{2+} influx (Meena *et al.*, 2019). Thus, it becomes increasingly evident that herbivory-induced $[Ca^{2+}]_{cyt}$ elevation involves multiple channels and pathways regulating local and long-distance $[Ca^{2+}]_{cyt}$ signals.

Nevertheless, some studies showed that conventional channels might not always be responsible for Ca^{2+} influx pathways. Thus, the involvement of other passive Ca^{2+} transport-mediating proteins, such as annexins, becomes an interesting possibility (Laohavisit & Davies, 2009, 2011; Davies, 2014; Ma *et al.*, 2019).

Annexins are found in eukaryotic organisms and form a diverse multigene superfamily of Ca^{2+} -dependent membrane-binding proteins that serve as targets for Ca^{2+} in most eukaryotic cells. In angiosperms, annexins are found in vegetative and generative organs (Laohavisit & Davies, 2011; Clark *et al.*, 2012). They are composed of motifs 60–70 amino acids long, repeated four times. The ability of annexins to conduct Ca^{2+} has become evident from *in vivo* and *in vitro* assays (Demidchik & Maathuis, 2007; Laohavisit *et al.*, 2009, 2012; Richards *et al.*, 2014; Ma *et al.*, 2019). Unlike conventional Ca^{2+} channels, which are

routed from the Golgi complex to reside in a specific membrane, annexin proteins are able to occupy multiple cellular locations simultaneously. This characteristic makes annexins capable of a fast-recruitment response that can be driven by localized stimulation of membrane regions and might be independent of vesicular delivery – reviewed by Laohavisit & Davies (2009, 2011) and Clark *et al.* (2012).

Among the eight annexins described to date in *Arabidopsis thaliana*, ANNEXIN1 (ANN1) is the best-studied one. It was initially detected in the cytosol of cells, and later in the plasma membrane, endoplasmic reticulum, vacuole, mitochondria, chloroplast, and in the cell wall (Laohavisit & Davies, 2011). *ANN1* overexpression has a protective effect on plant survival under drought conditions, whereas lack of expression increases stress sensitivity (Konopka-Postupolska *et al.*, 2009). Studies on *Arabidopsis* roots have correlated its localization in the plasma membrane with the presence of a Ca^{2+} conductance, which is activated by voltage hyperpolarization and extracellular hydroxyl radicals and is involved in the elongation of root hair cells (Foreman *et al.*, 2003; Laohavisit *et al.*, 2012). The *Arabidopsis ann1* knockout mutant lacks this Ca^{2+} -channel-like activity in the plasma membranes of root epidermal cells and root hairs. Furthermore, *ann1* mutants also have shorter roots compared to wild-type (Columbia-0, Col-0) plants (Laohavisit *et al.*, 2012). More recent studies have shown that ANN1 is involved in root and seedling $[Ca^{2+}]_{cyt}$ elevation in response to hydrogen peroxide (Richards *et al.*, 2014; Zhao *et al.*, 2019).

As ANN1 is firmly implicated in $[Ca^{2+}]_{cyt}$ elevation and this occurs during insect feeding, our aim was to elucidate a putative role for this annexin in plant responses to herbivory-related cues. Therefore, we performed a set of assays to characterize *ann1* mutants. The effect of the lack of ANN1 was analyzed by observing larval growth of the crop-pest moth *Spodoptera littoralis* on two different *ann1* knockout and two ANN1-overexpressing lines. Further, the plant's response to mechanical injuries (i.e. mechanical wounding with and without the addition of larval OS) and after lesions caused by *S. littoralis* feeding on leaves was investigated. We found a role for ANN1 in *Arabidopsis* for both local and systemic defense responses against *S. littoralis* attack by mediating $[Ca^{2+}]_{cyt}$ elevations, jasmonate level, and defense-related gene expression. This study contributes to our understanding of the molecular identity of Ca^{2+} channels involved in the plant response to wounding and herbivory.

Materials and Methods

Plant growth and treatment

Arabidopsis thaliana Col-0 wild-type, *ANN1* knockout mutant lines *ann1-1* (SALK_015426 originally characterized by Lee *et al.* (2004) and *ann1-2* (WiscDsLox477–480P11), and *ANN1*-overexpressing lines *ANN1-OE10* and *ANN1-OE12* (Konopka-Postupolska *et al.*, 2009) were used. For $[Ca^{2+}]_{cyt}$ measurements, Col-0 and *ann1-1* were transformed by floral dip to express cytosolic (apo)aequorin driven by the 35S promoter. Floral dip employed *Agrobacterium tumefaciens* GV3101 pNEW 35S::AEQ

pGreen vector containing the 35S::AEQ insert cut from pMAQ (Knight *et al.*, 1991). All plants used in aequorin assays were at least three generations posttransformation.

Plants were kept in short-day conditions after stratification for 2 d at 4°C. Four to five-week-old plants grown in 10 cm round pots were used for all experiments. The growth chamber was adjusted to 50–60% relative humidity and 21°C with a 10 h : 14 h, light : dark photoperiod and a light intensity of 100 $\mu\text{mol m}^{-2} \text{s}^{-1}$. For experiments investigating the systemic response and translocation of metabolites, the leaves of each plant were counted according to their age (Dengler, 2006; Farmer *et al.*, 2013; Kiep *et al.*, 2015).

MecWorm (Mithöfer *et al.*, 2005) treatment was used for mechanical wounding of the plant with punches every 5 s (12 punches per minute) on treated leaf 8. To investigate the systemic response upon treatment of leaf 8, the local and systemic leaves 5, 8, 9 and 13 were analyzed. Untreated plants were used as control and had the same growth and handling conditions as the treated ones.

To study the mechanical-wounding-induced systemic response, wounds were generated on leaf 8 with a pattern wheel (six vertical movements on each side of the midrib), and 20 μl of water (MW + W: mechanical wounding + water) or of *S. littoralis* OS, diluted 1 : 1 in water (MW + OS) was applied to the wounds (Vadassery *et al.*, 2012a). Treated plants were kept in the growth chamber with a cover to prevent evaporation. Samples of leaf 8 and selected systemic leaves were harvested in liquid N₂ and kept at –80°C till further analysis.

Analysis of [Ca²⁺]_{cyt} elevations

For the analysis of [Ca²⁺]_{cyt} in whole plants, leaves of 4-wk-old Arabidopsis rosettes were numbered according to their phyllotactic sequence (Dengler, 2006). The day before the experiment, plants were sprayed with 10 μM coelenterazine in 0.01% (v/v) Tween 20 and incubated in the dark for 16 h for aequorin reconstitution. Aequorin imaging was performed according to Kiep *et al.* (2015) using a high-resolution photon-counting camera system (HRPCS218; Photek, St Leonards-on-Sea, UK) comprising an intensified CCD camera (ICCD218; Photek) and a camera controller (HRPCS4; Photek). The camera was mounted on a darkbox (DB-2; Photek). Signal acquisition and processing were performed with the IFS32 software (Photek). Photons were captured in photon-counting mode with a 200 ms frame rate, and cumulative images were integrated offline after the experiments as indicated in the figure legends. At the end of each treatment, the rosettes were flooded with 40 ml discharge solution (1 M calcium chloride, 10% (v/v) ethanol) to achieve a complete discharge of aequorin to enable calibration of the data obtained and determine the cytosolic Ca²⁺ concentration according to Knight *et al.* (1996). The identical regions of interest (ROIs) found in treatment and discharge images were identified, and the average signal intensity in the ROIs at a given time point, as well as the cumulative counts in the ROIs, were determined by using the IFS32 software. The wounding treatment consisted of mechanically wounding the midrib of leaf 8 with a pattern wheel and

adding 20 μl of water (MW + W) or 20 μl of OS (MW + OS) across all holes of the mechanically wounded leaf.

Insect material and feeding assays

Larvae of the generalist herbivore *S. littoralis* were hatched and reared on artificial diet at 23–25°C with 10 h : 14 h, light : dark cycles (Bergomaz & Boppré, 1986). The OS was collected from *S. littoralis* larvae fed on *Arabidopsis* Col-0 plants and stored on ice. The OS was centrifuged at 10 000 g and 4°C to remove residual plant tissue pieces and then diluted with water (1 : 1) as previously described (Vadassery *et al.*, 2012a).

For short-term feeding assays, third instar *S. littoralis* larvae were used after being kept separately without food overnight. This treatment ensured an immediate start of feeding after placement on the plant. The locally fed leaves were collected in liquid N₂ after the indicated time points and kept at –80°C until further analysis.

For local larval feeding assays, overnight-starved third-instar larvae were placed on leaf 8 for direct feeding. Each plant received one larva. After feeding on *c.* 40% of the leaf, which took between 5 and 10 min, the larva was removed. After 90 min, the local and systemic leaves were harvested for phytohormone extraction and gene expression analysis.

For 1 wk feeding assays, 30 first-instar larvae were placed on each of 10 Col-0, *ann1* and *ANN1-OE* plants (three larvae per plant). To achieve similar starting conditions, all larvae determined for one plant genotype were pooled and weighed before the experiment. The minimal starting weight of 30 larvae was set to 60 mg. After 1 wk, the weight of all larvae found again was recorded separately. Owing to a limited number of first instar larvae available, the experiment was carried out several times and the weight data of each genotype were combined.

Gene expression

Total RNA was extracted from frozen material (*c.* 50–100 mg) using the Trizol method according to the manufacturer's protocol (Thermo Fisher, Darmstadt, Germany). Genomic DNA in total RNA samples was removed using the Turbo DNase-free kit (Ambion, Thermo Fisher) according to the manufacturer's protocol. The integrity and amount of RNA were monitored by agarose gel electrophoresis and spectrophotometric quantification, respectively. Complementary DNAs were synthesized using the GeneAmp Core PCR RNA Kit (Applied Biosystems) according to the manufacturer's instructions. The pair of primers specific for *ANN1* (AT1G35720: forward 5'-ATGGCGACTCTTAA GGTTCCTGAT-3' according to Clark *et al.* (2001) and reverse 5'-GCCTGATGACTTTCCTCTGTTTCAG-3') was used, producing a product size of 151 bp. For *VEGETATIVE STORAGE PROTEIN2 (VSP2; AT5G24770)* we used forward 5'-ACGACT CCAAACCGTGTGCAA-3' and reverse 5'-CGGGTTCGGT CTTCTCTGTTCCGT-3' (Vadassery *et al.*, 2012b), and for *JASMONATE-ZIM-DOMAIN PROTEIN 10 (JAZ10; AT5G13220)* we used forward 5'-TCGAGAAGCGCAAGGA GAGATTAGT-3' and reverse 5'-AGCAACGACGAAGAAGG

CTTCAA-3' (Scholz *et al.*, 2014). Quantitative reverse transcription PCR was performed on a CFX96 Real Time System (Bio-Rad). Brilliant II QPCR SYBR green Mix (Agilent, Böblingen, Germany) was used to monitor the synthesis of double-stranded DNA. Each biological sample was analyzed in technical triplicates. The cycle protocol consisted of 95°C for 10 min, followed by 40 cycles of 95°C for 15 s and 60°C for 1 min, and ended by a dissociation curve determined between 60°C and 95°C. The specificity of PCR amplifications was evaluated by the presence of a single peak in denaturation curves and by visualization of simple amplification products of expected size in ethidium bromide gel electrophoresis. The primer efficiencies were calculated using LINREGPCR (v.11.0, (Ruijter *et al.*, 2009)). The mean relative expression of the gene was calculated according to Pfaffl (2001), using $\Delta\Delta C_t$ with the *ribosomal protein S18* gene (AT1G34030) as reference (Scholz *et al.*, 2014) (*RPS18*, forward 5'-GTCTCCAATGCCCTTGACAT-3'; reverse 5'-TCTTTCC TCTGCGACCAGTT-3').

Extraction and quantification of phytohormones

A total of 250 mg of leaf material was used for phytohormone analyses. The extraction procedure and the determination of JA and JA-Ile were performed as previously described (Vadassery *et al.*, 2012a) with some modifications. An API5000 triple quadrupole mass spectrometer (AB Sciex, Darmstadt, Germany) was used for detection (Heyer *et al.*, 2018a). Moreover, in this study, a different mixture of labeled jasmonates was used as internal standard. Instead of 15 ng of JA- $^{13}C_6$ -conjugate used in the previous study, 12 ng of D₆-JA-Ile (HPC Standards GmbH, Cunnorsdorf, Germany) was used. In addition, the 60 ng of 9,10-D₂-9,10-dihydrojasmonic acid was replaced by 60 ng of D₆-JA (HPC Standards GmbH) as previously reported by Scholz *et al.* (2017).

Statistics

To ensure reproducibility, all experiments were repeated with independent biological replicates. The exact number of replicates is indicated in the particular figure legends. For statistical analyses, one or two-way ANOVA followed by *post hoc* (Student–Newman–Keuls; Šidák; Tukey) tests or Student's *t*-tests were used as indicated in the figure legends. Different letters indicate significant differences between treatments or leaves. GraphPad PRISM 6 and ORIGINPRO 9.3 software were used for data analysis and graph composition.

Results

ANNEXIN1 is induced by mechanical wounding and herbivory

Initially, in order to learn whether ANN1 was involved in the plant's defense response against wounding and herbivory, we implemented different wounding-related treatments on Col-0 and *ann1-1* knockout plants and then analyzed ANN1 expression

levels. Different treatments were performed on leaf number 8: first, mechanical wounding (MW) with a pattern wheel and applying water (MW + W) or OS of *S. littoralis* (MW + OS) simulating herbivory; and second, direct feeding of *S. littoralis* larvae on one leaf. Col-0 plants showed a high accumulation of ANN1 transcripts after all treatments, whereas in the *ann1-1* mutant no induction of ANN1 was detected (Fig. 1). No significant difference between the different treatments was detected (Fig. 1a), suggesting a universal response of ANN1 expression in response to mechanical wounding, with or without the presence of larval OS, and to herbivore attack.

Local and systemic calcium signaling is affected in *ann1* plants

Since, on the one hand, a $[Ca^{2+}]_{cyt}$ signal precedes jasmonate accumulation and subsequent defense-related responses upon wounding and herbivory (Fisahn *et al.*, 2004; Maffei *et al.*, 2007; Bricchi *et al.*, 2010; Toyota *et al.*, 2018; Meena *et al.*, 2019) and, on the other hand, annexins are components of $[Ca^{2+}]_{cyt}$ signal generation, we aimed to investigate how the elevation of $[Ca^{2+}]_{cyt}$ upon stress is influenced by the presence or absence of the ANN1 protein. Therefore, we used Col-0 and *ann1-1* plants, both containing the $[Ca^{2+}]_{cyt}$ reporter (apo)aequorin. The $[Ca^{2+}]_{cyt}$ elevation was analyzed in whole-plant rosettes according to Kiep *et al.* (2015). Leaf 8 was wounded with a pattern wheel before adding




Fig. 1 Response of ANNEXIN1 (*ANN1*) in *Arabidopsis thaliana* after treatment with different stresses. Levels of ANN1 transcripts (\pm SE) were determined on (a) the wild-type (Col-0) and (b) the *ann1-1* mutant genotype after injuring leaf 8 with a pattern wheel, applying either water (MW + W) or oral secretion (MW + OS) to the wounds, or promoting *Spodoptera littoralis* larvae feeding (Larvae, third instar) feeding on leaf 8 until c. 40% of the leaf was eaten. Per genotype and treatment, nine replicates ($n = 9$) were done. All plants were incubated for 90 min before sampling leaves. Leaves of untreated plants were used as controls (Ctrl). Differences between treatments within the same genotype were analyzed using two-way ANOVA (Student–Newman–Keuls *post hoc* test); significant differences are indicated by different letters ($P < 0.05$); ns, not significant

20 μl water or OS (1 : 1 diluted) to the wounds. We observed an immediate and monophasic local elevation of $[\text{Ca}^{2+}]_{\text{cyt}}$ with a peak after *c.* 30–45 s, and a return to background levels after *c.* 4 min (Fig. 2; Supporting Information Fig. S1; Videos S1–S4). The local response was similar to those described in previous studies, but presenting a slightly faster $[\text{Ca}^{2+}]_{\text{cyt}}$ peak response (cf. 1 min: Verrillo *et al.*, 2014; Kiep *et al.*, 2015). In the case of MW + OS treatment, the Col-0 $[\text{Ca}^{2+}]_{\text{cyt}}$ elevation lasted longer than with the MW + W (water) treatment. Although slightly weaker than in the Col-0 plants, the local $[\text{Ca}^{2+}]_{\text{cyt}}$ response was clearly detectable in the wounded *ann1-1* leaf and moved into the petiole. Strikingly, neither water nor OS induced a systemic $[\text{Ca}^{2+}]_{\text{cyt}}$ response in the *ann1-1* genotype (Figs 2, S1). By contrast, upon MW + W treatment, a systemic $[\text{Ca}^{2+}]_{\text{cyt}}$ response (also monophasic) was observed in Col-0 plants (systemic leaf 5), which started after 4 min and reached a maximum at 4.5 min before decreasing to background levels (Fig. S1). Upon MW + OS treatment, the systemic response was more pronounced. Here, leaves 5, 6, 7, 10 and 11 responded in Col-0 (Fig. 2c). This response was highly significantly different when compared with *ann1-1*. It was possible to observe a $[\text{Ca}^{2+}]_{\text{cyt}}$ wave running from the treated leaf to the connected leaves in Col-0 (Videos S1, S3), whereas only the local response was detected in treated *ann1-1* plants (Videos S2, S4). Therefore, ANN1 seems to be an important player in the systemic $[\text{Ca}^{2+}]_{\text{cyt}}$ wave upon wounding and herbivory challenge.

ann1 plants are more susceptible to herbivore feeding, whereas ANN1-overexpressing plants are more resistant

To further evaluate the impact of ANN1 on the defense against chewing herbivores, we carried out feeding assays using first instar *S. littoralis* larvae on Col-0 wild-type plants and two different

knockout lines (*ann1-1*, *ann1-2*) and two different ANN1 overexpressor lines (*ANN1-OE12*, *ANN1-OE10*). We performed two independent sets of experiments, each with Col-0 as wild-type control, one knockout, and one overexpressor line. Our results show that larvae feeding on *ann1* plants gained significantly more weight than wild-type-fed larvae (Fig. 3). This effect was also observed when we evaluated the larval growth on the APOAEQUORIN-containing *ann1* mutant (*ann1-1/AEQ*) (Fig. S2). The opposite happened when the larvae were feeding on the ANN1-OE plant lines, where they gained significantly less weight than those feeding on Col-0 plants (Fig. 3).

Spodoptera littoralis feeding-induced jasmonate accumulation is affected in ANN1 mutant plants

Jasmonates are rapidly induced upon wounding and herbivory. To test if these were differentially induced in *ann1* or ANN1-OE10 lines compared with the Col-0 wild-type, we analyzed jasmonate concentrations after herbivore feeding. In independent short-term (30, 90 min) feeding assays, the levels of both JA and JA-Ile increased significantly in the treated leaves of all genotypes at both time points (Fig. 4). Even fed leaves of *ann1-1* and *ann1-2* plants still accumulated both JA and JA-Ile; however, their levels were significantly lower than in Col-0 plants (Fig. 4). By contrast, in the overexpressing line ANN1-OE10 the level of JA-Ile was significantly higher than in Col-0, although the JA level was not (Fig. 4c,d).

Systemic transcriptional responses to *S. littoralis* attack are impaired in *ann1-1* plants

In order to elucidate whether ANN1 is also involved in systemic defense, the effect of herbivory on local and systemic defense




Fig. 2 Cytosolic free calcium ($[\text{Ca}^{2+}]_{\text{cyt}}$) response upon mechanical wounding (MW) and oral secretion (OS) in Col-0 and annexin1 mutant (*ann1-1*). (a) Cumulative images over 10 min of local and systemic $[\text{Ca}^{2+}]_{\text{cyt}}$ signals in 4-wk-old whole *Arabidopsis thaliana* Col-0 (top) and *ann1-1* (bottom) rosettes in response to MW + OS applied to leaf 8 (left) and after discharge (right). Discharge was used to determine L_{total} to calculate the $[\text{Ca}^{2+}]_{\text{cyt}}$ in individual leaves. (b) Kinetics of a representative $[\text{Ca}^{2+}]_{\text{cyt}}$ response in individual leaves in Col-0 and *ann1-1* after treatment. (c) $[\text{Ca}^{2+}]_{\text{cyt}}$ response (mean \pm SE) in individual leaves in Col-0 and *ann1-1* after treatment. Per genotype and treatment, nine replicates ($n = 9$) were done. cul, cumulative; dis, discharge. Two-way ANOVA (Sidak *post hoc* test; ***, $P < 0.001$).




Fig. 3 Weight of *Spodoptera littoralis* larvae after feeding on *Arabidopsis thaliana* Col-0 plants, two annexin1 (*ann1*) mutants, and two *ANN1*-overexpression lines. Thirty first-instar larvae of *S. littoralis* were preweighed, and three larvae were placed on each plant. In each independent experiment, 10 plants per genotype were used. Larva weight was measured individually after 7 d of feeding. Each set of experiments was independently repeated: (a) $n = 3$; (b) $n = 4$. The combined total number of larvae n recovered after 1 wk is indicated. The box indicates the middle 50% of the data points; the black line within the box is the median. Whiskers are defined as 1.5-fold interquartile range; dots represent outliers. Statistical differences between the genotypes after feeding were analyzed using one-way ANOVA (Tukey's *post hoc* test) and indicated by different letters ($P < 0.001$).

responses was analyzed in parallel, focusing on the full knockout line *ann1-1* (Fig. 1b). We performed an experiment in which *S. littoralis* was allowed to feed on one defined local leaf (leaf 8) followed by leaf sampling after 90 min – leaf numbering according to Dengler (2006), Farmer *et al.* (2013), and Kiep *et al.* (2015). In addition to the treated leaf 8, unwounded systemic

leaves 5 and 13 (vascularly connected to leaf 8) and leaf 9 (unconnected to leaf 8) were sampled and analyzed for the expression of two jasmonate-responsive genes, *VSP2* and *JAZ10*. Compared with the nontreated Col-0 plants, *VSP2* was induced 10-fold in the local leaf 8 and even more strongly in the directly vascularly connected leaf 13, whereas no change in expression was




Fig. 4 Local accumulation of jasmonates in *Arabidopsis thaliana* leaves after short-time feeding assay with *Spodoptera littoralis*. (a, c) Jasmonic acid (JA) and (b, d) (+)-7-*iso*-jasmonoyl-L-isoleucine (JA-Ile) levels (mean \pm SE) from *A. thaliana* leaves were analyzed after 30 and 90 min of feeding from a single third instar *S. littoralis* larva. Phytohormones were determined only from fed leaves. Five to eight replicates ($n = 5-8$) were done per time point. Leaves of untreated plants were used as controls. The differences of JA and JA-Ile levels between time points and genotypes were analyzed using two-way ANOVA (Sidak *post hoc* test); significant differences are indicated by different letters ($P < 0.005$).

detectable in the *ann1-1* mutant (Fig. 5a). Also, *JAZ10* was significantly induced in the treated leaf 8 in Col-0 as well as in all systemic leaves, again with the highest expression in leaf 13. In addition, in *ann1-1* plants, *JAZ10* showed a significant induction in systemic leaves 5 and 13 (Fig. 5b).

Herbivory-like feeding-induced systemic jasmonate accumulation is abolished in *ann1-1* plants

Considering that ANN1 is involved in jasmonate-related defense induction in systemic leaves, we wanted to study whether mechanical wounding alone employs ANN1 or if a chemical signal of the OS of the larvae is necessary. Thus, we tested the systemic jasmonate response after local leaf wounding with a pattern wheel followed by either water or OS application to the small wounds. The results show a clear local jasmonate response in Col-0 and in *ann1-1* plants upon wounding and water (Fig. 6). This local response was significantly higher when wounded sites were treated with OS. Interestingly, only OS treatment induced a systemic response in the vascularly connected leaves 5 and 13 in Col-0, represented by a strong increase of JA and JA-Ile. This was not the case in *ann1-1* plants, where the systemic effect was completely absent (Fig. 6). These results were supported by other measured jasmonates. The biosynthetic precursor *cis*-12-oxo-phytodienoic acid and the catabolite hydroxy-JA both showed very similar results to those found for JA and JA-Ile (Fig. S3).

Previous work has shown that pattern-wheel wounding does not fully represent the insect feeding-like wounding such as, for example, the use of MecWorm does (Mithöfer *et al.*, 2005).

Therefore, MecWorm-mediated wounding was applied in an additional experiment. Based on the studies of Heyer *et al.* (2018b), we wounded leaf 8, including the midrib. As shown in Fig. 7, the continuous mechanical wounding of leaf 8 inflicted by MecWorm significantly elevated the local levels of JA-Ile, which we focused on as the bioactive form of the jasmonates, in both Col-0 and *ann1-1* plants. Strikingly, in contrast to Col-0, in *ann1-1* plants, local accumulation of jasmonates in leaf 8 was not accompanied by a comparable systemic increase of jasmonates. Moreover, a highly significant difference in JA-Ile accumulation in directly connected leaf 13 was observed when Col-0 and *ann1-1* plants were compared (Fig. 7).

Discussion

Various studies have demonstrated that a rapid, early and transient increase of $[Ca^{2+}]_{cyt}$ is involved and essential for the successful induction and regulation of jasmonate accumulation and further downstream plant defense strategies upon wounding and insect herbivory (Fisahn *et al.*, 2004; Maffei *et al.*, 2004, 2007; Arimura *et al.*, 2008, 2011; Scholz *et al.*, 2014; Toyota *et al.*, 2018; Yan *et al.*, 2018; Kumari *et al.*, 2019; Meena *et al.*, 2019). Such stress-induced $[Ca^{2+}]_{cyt}$ elevation occurs both locally and systemically (Kiep *et al.*, 2015). There are various channels that have been shown to be involved in wounding or herbivory-related Ca^{2+} influx into the cytosol such as the TPC1 (Kiep *et al.*, 2015; Vincent *et al.*, 2017), glutamate receptor-like channels (GLRs; Mousavi *et al.*, 2013; Toyota *et al.*, 2018), and cyclic nucleotide-gated channel 19 (CNGC19; Meena *et al.*, 2019). However,




Fig. 5 Local and systemic transcriptional responses to *Spodoptera littoralis* feeding. (a) Levels of *VEGETATIVE STORAGE PROTEIN2* (*VSP2*) and (b) *JASMONATE-ZIM-DOMAIN PROTEIN 10* (*JAZ10*) transcripts were determined in *Arabidopsis thaliana* Col-0 and annexin1 (*ann1-1*) genotypes in different leaves after local *S. littoralis* (Larvae) feeding for 90 min on leaf 8 (t). Treated leaf 8 and untreated leaves 5, 9 and 13 were analyzed (mean \pm SE). Per genotype and treatment, nine replicates ($n = 9$) were done. Leaves of untreated plants were used as controls. *RPS18* was used as a reference gene to normalize the data. Statistical differences between treatments were analyzed using two-way ANOVA (Student–Newman–Keuls *post hoc* test); significant differences are indicated by different letters ($P < 0.05$).




Fig. 6 Local and systemic accumulation of jasmonates after mechanical wounding (MW) and oral secretion (OS) treatment. Levels of (a) jasmonic acid (JA) and (b) (+)-7-*iso*-jasmonoyl-L-isoleucine (JA-Ile) were determined in *Arabidopsis thaliana* Col-0 and annexin1 (*ann1-1*) genotypes in different leaves 90 min after wounding of leaf 8 with a pattern wheel (MW) and applying water (MW + W) or oral secretion (MW + OS) on the wounds (mean \pm SE). Per genotype and treatment, seven replicates ($n = 7$) were done. Untreated leaves 5, 9 and 13 and treated leaf 8 (t) were analyzed. Leaves of untreated plants were used as controls. Statistical differences found in leaves between treatments, or between genotypes were analyzed using two-way ANOVA (*, $P < 0.05$; **, $P < 0.005$; ***, $P < 0.001$; Student–Newman–Keuls test). No indication, no significant difference.




Fig. 7 Accumulation of (+)-7-*iso*-jasmonoyl-L-isoleucine (JA-Ile) in leaves of *Arabidopsis thaliana* after mechanical wounding with MecWorm. JA-Ile levels were analyzed in Col-0 and annexin1 (*ann1-1*) genotypes in different leaves after mechanical wounding for 90 min, including the midrib. In treated plants, leaf 8 (t) was subjected to mechanical damage; untreated leaves 5, 9 and 13 and treated leaf 8 were analyzed (mean \pm SE). Per genotype and treatment, six to eight replicates ($n = 6–8$) were done. Leaves of untreated plants were used as controls. Statistical differences between leaves of control and treated plants on the same genotype and between Col-0 and *ann1-1* leaves upon MecWorm treatment were analyzed using an unpaired Student's *t*-test (*, $P < 0.05$; **, $P < 0.005$). No indication, no significant difference.

Ca^{2+} influx may not be mediated only by conventional channels – additional, unconventional ones such as annexins might also contribute (Laohavisit & Davies, 2011; Laohavisit *et al.*, 2012; Davies, 2014; Ma *et al.*, 2019). Diverse studies in plants have gathered evidence of annexins' ability to influence Ca^{2+} transport. A growing body of data suggests that annexins play a role in plant response to nematode parasitism. Some cyst-secreted effectors are annexin-like and able to affect plant defense possibly by mimicking the endogenous annexin functions and impairing H_2O_2 -induced $[\text{Ca}^{2+}]_{\text{cyt}}$ transients (Patel *et al.*, 2010; Zhao *et al.*, 2019). Moreover, it was shown that compared with wild-type, overexpression of *ANN1* and *ANN4* decreases susceptibility against *Meloidogyne incognita* nematode infection of roots while the *ann1* and *ann4* lines were more susceptible (Zhao *et al.*, 2019). Our study demonstrates a role for *ANN1* in systemic leaf defense responses against herbivore attack and mechanical wounding.

ANNEXIN1 is induced upon insect herbivory

Biotic interaction has been shown to influence annexin transcription in crops such as alfalfa (*Medicago sativa*), Indian mustard (*Brassica juncea*), tomato (*Solanum lycopersicum*), and wheat (*Triticum aestivum*) (Kovács *et al.*, 1998; Jami *et al.*, 2009; Lu *et al.*, 2012; Xu *et al.*, 2016). It was shown in multiple studies that *Arabidopsis ANN1* gene expression is influenced by diverse environmental signals (Konopka-Postupolska *et al.*, 2009; Clark *et al.*, 2010; Guelette *et al.*, 2012). Here, we demonstrate the expression response of *Arabidopsis ANN1* to herbivory. The assays were designed to understand the attack in a mechanistic and holistic way, which we achieved by dissecting the insect

attack into different modules of stress: mechanical wounding alone (MW + W), mechanical wounding plus OS (MW + OS), or the complex stress of larval feeding (Fig. 1). The high expression of *ANN1* upon the different but complementary stresses showed that *ANN1* transcription activation is triggered quickly (here, after 90 min) by insect-feeding-related damage, but also by wounding alone. The latter confirms earlier results showing *ANN1* induction 24 and 48 h after wounding (Konopka-Postupska *et al.*, 2009).

ANNEXIN1 is involved in the defense against herbivores

As with other cellular components that are involved in $[Ca^{2+}]_{cyt}$ signaling (Vadassery *et al.*, 2012b; Scholz *et al.*, 2014; Meena *et al.*, 2019), we show here that ANN1 is also an important player in the regulation of insect-feeding-induced defense. Using two different knockout and two different overexpression lines, we showed that *S. littoralis* larvae feeding on *ann1* plants gained substantially more weight (in total, +27.6%), whereas those feeding on ANN1-OE plants were much smaller (−26.9%) than larvae feeding on Col-0 plants (Figs 3, S2). Thus, ANN1 is a positive regulator in herbivory-induced defense in Arabidopsis and contributes to resistance against *S. littoralis*, comparable to CNGC19 and GLRs.

To gain further insight into the putative role of ANN1 in systemic defense-related signaling, we performed a series of experiments in which a defined local leaf was treated and systemic leaves were analyzed (Mousavi *et al.*, 2013; Kiep *et al.*, 2015). Using (apo)aequorin-expressing *ann1-1* and wild-type plants, we demonstrated that ANN1 is indispensable for the systemic $[Ca^{2+}]_{cyt}$ response; this is comparable to what has been found previously for TPC1 and GLRs (Kiep *et al.*, 2015; Toyota *et al.*, 2018). By contrast, the local response was slightly but not significantly reduced in the *ann1-1* mutant (Figs 2, S1). However, compared with other studies (Nguyen *et al.*, 2018; Toyota *et al.*, 2018), the systemic $[Ca^{2+}]_{cyt}$ response was found to be weaker. This can be explained by different experimental conditions; first, in the other studies, the GCaMP3 fluorescent-protein-based $[Ca^{2+}]$ sensor was used, a highly sensitive calcium fluorescence reporter, whereas we used bioluminescent aequorin. Second, their mode of wounding was much harsher. Whereas Nguyen *et al.* (2018) destroyed half of the leaf tissue, we used a pattern wheel that caused only a few small holes in the tissue. This supports the view that the intensity of wounding correlates with the intensity of the response (Nguyen *et al.*, 2018).

As a readout for the defense response, the expression of jasmonate-responsive genes *VSP2* and *JAZ10* was examined. Upon larval feeding, a local increase was detected in Col-0 for both genes, as well as a strong increase in the vascularly connected leaf 13 that was even higher than the local response (Fig. 5). Such a strong systemic increase was not found for either *VSP2* or *JAZ10* in *ann1-1* plants, suggesting that the feeding-related signals necessary to induce the systemic gene activation do not reach the distal leaves. The fact that in this and other experiments leaf 13 and (to a lesser extent) leaf 5 respond more strongly can be explained by the direct (leaf 13) and indirect (leaf 5) vascular connections to leaf 8 (Dengler, 2006).

As $[Ca^{2+}]_{cyt}$ elevations initiate Ca^{2+} signaling (Kudla *et al.*, 2010; Mithöfer & Boland, 2012; Scholz *et al.*, 2014; Vadassery *et al.*, 2014) and precede downstream signals, such as phytohormone accumulation, we further investigated the concentrations of the jasmonates JA and JA-Ile at two time points (30 and 90 min). These jasmonates were found to be strongly induced locally in both Col-0 and *ann1* upon herbivore attack; but accumulation was significantly lower in *ann1* plants, whereas in the *ANN1-OE10* at least the JA-Ile level was significantly higher (Fig. 4). This is in accordance with the finding that $[Ca^{2+}]_{cyt}$ signals in *ann1* plants are affected in local leaves to a certain extent and suggests that a full induction of jasmonates in response to insect herbivores is not possible in those plants. Strikingly, the induction of *ANN1* upon herbivory or wounding (Fig. 1) is high if compared with downstream responses such as jasmonate or gene induction. Very likely, the reason behind this finding is that gene expression does not always reflect the corresponding protein expression. Lee *et al.* (2004) already noted that *ann1-1* transcript level does not necessarily correspond to the ANN1 protein. Besides using larvae, we further evaluated the phytohormonal responses in local and systemic leaves of plants treated with mechanical wounding. We chose this approach because such experiments can be better standardized than exposure to larvae that might feed or not, particularly when only a short time period is investigated. Pattern-wheel-mediated mechanical wounding, as well as wounding by MecWorm, supported the finding of local jasmonate induction in wild-type and mutant (Figs 6, 7). Strikingly, after MW + OS treatment, the response of JA-Ile in *ann1-1* plants was significantly reduced compared with wild-type (Fig. 6). In addition, an elevation of jasmonates was also observed in systemic leaves of OS-treated plants in wild-type but not in *ann1-1* mutant plants. This further indicates that ANN1 is involved in systemic $[Ca^{2+}]_{cyt}$ -dependent jasmonate elevation. This notion was supported by results obtained by mechanical wounding alone, using MecWorm treatment. In *ann1-1* plants, leaf 13 showed significantly lower accumulation of JA-Ile than in Col-0 plants (Fig. 7), suggesting that ANN1 might not only be involved in systemic OS-specific signaling but also in systemic wound-induced jasmonate accumulation.

Conflating the data obtained, we propose that ANN1 is a positive factor of the $[Ca^{2+}]_{cyt}$ -dependent systemic defense response against wounding and larval feeding. Thus, the absence of this unconventional Ca^{2+} channel causes an impaired systemic response. ANN1 can exist as an integral plasma membrane protein (Alexandersson *et al.*, 2004; Marmagne *et al.*, 2007). Nevertheless, as small amphipathic proteins, annexins are distributed throughout cells and can be transported within the plant via the phloem (Guelette *et al.*, 2012). It is possible that annexins may be recruited directly to membranes, independently of vesicle delivery, to operate in stimulus-specific signaling (Laohavisit & Davies, 2011; Laohavisit *et al.*, 2012; Davies, 2014; Espinoza *et al.*, 2017). Therefore, a plant that contains the conventional and unconventional Ca^{2+} channels should be able to recruit annexins to the tissues under stress when needed, whereas the *ann1* genotype can only launch reduced $[Ca^{2+}]_{cyt}$ -mediated defense responses. It should be kept in mind that annexins might

also act via the regulation of channel activities, for example by selective channel delivery to or retraction from membranes, in a similar way as the KAT1 plasma membrane potassium ion (K^+) channel is cycled during abscisic acid induced stomatal closure (Sutter *et al.*, 2007). However, the observed effect of abolishing systemic $[Ca^{2+}]_{cyt}$ -induced responses in *ann1* plants may be specific for certain stresses, such as herbivory, and is not necessarily involved in all types of biotic and abiotic stress responses. An example is a study showing that ANN1 was not necessary for systemic signaling and development of acquired resistance in uninfected leaves during challenge with avirulent bacteria *Pseudomonas syringae* pv tomato (Carella *et al.*, 2016).

The success of the plant's defense response against stresses does not depend on one single signaling component alone. Instead, it is a coordinated local and systemic communication between cells and distant organs. For that, various signals, such as ROS, hydraulic pressure, as well as electropotential waves, $[Ca^{2+}]_{cyt}$ and others, are employed in a tightly linked manner (Foreman *et al.*, 2003; Zimmermann *et al.*, 2009; Maischak *et al.*, 2010; Farmer *et al.*, 2013; Mousavi *et al.*, 2013; Davies, 2014; Seybold *et al.*, 2014; Ranjan *et al.*, 2015; Peiter, 2016; Alonso *et al.*, 2019; Gully *et al.*, 2019; Saijo & Loo 2019). In previous studies using epidermal root tissue, hydroxyl-radical-activated plasma membrane conductance of Ca^{2+} and K^+ were absent in the *ann1-1* mutant (Laohavisit *et al.*, 2012). Expression and protein levels of Arabidopsis ANN1 also correlated with the occurrence of the radical-activated plasma membrane Ca^{2+} conductance in the root epidermis and at the apex of root hairs (Clark *et al.*, 2001; Dinnyen *et al.*, 2008). These results strongly suggest that ANN1 is very likely a Ca^{2+} -permeable protein in Arabidopsis and might provide a molecular link between ROS and $[Ca^{2+}]_{cyt}$ in the systemic defense-related signaling in plants. Further studies will address this hypothesis.

In conclusion, we investigated the role of ANN1 in local and systemic plant defense against wounding and herbivorous insects in Arabidopsis. Plant tissue wounding and cell disruption caused by feeding insects strongly induced ANN1 expression, demonstrating that it is part of the rapid defense response against invertebrate pests; neither jasmonates nor defense-related genes were upregulated systemically in *ann1* mutants. ANN1 mediates plant defense, affecting larval growth, and is crucial for the induction of signaling upon herbivory within the whole plant. ANN1 is an important part of systemic $[Ca^{2+}]_{cyt}$ signaling, thereby connecting $[Ca^{2+}]_{cyt}$ to subsequent downstream signals and defense responses against herbivores.

Acknowledgements

We thank A. Lehr, T. Peiter-Volk, and S. Janster for assistance in the laboratory, and the MPI-CE glasshouse team for growing plants. We additionally thank Wilhelm Boland, the Max Planck Society, and Celia R. Carlini for support. This work was further supported in part by the German Academic Exchange Service (DAAD; PPP project ID 57142556) and the Brazilian agency Coordenadoria de Aperfeiçoamento de Pessoal de Nível Superior (CAPES), Program PROBRAL 407/2016, grant 23038.006810/

2014-51 (JM), the International Max Planck Research School (AKM), the BBSRC (grant BB/K009869/1: KAW and JMD), the National Science Centre, Poland (2015/19/B/NZ3/01476: DK-P), and the DFG (CRC1127: SSS, RO; PE1500/7-1: EP). The authors declare that the research was conducted in the absence of any commercial or financial relationships that could be construed as a potential conflict of interest.

Author contributions

JM and AM conceived the project and research plans; JM, AKM, and MR performed the experiments; JM, AKM, SSS, RO and AM designed the experiments and analyzed the data; EP provided and assisted with the equipment for Ca analyses; DK-P, JR, JMD and KAW generated and provided seed material; JM, AKM, SSS and AM wrote the article with contributions from all authors.

ORCID

Julia M. Davies  <https://orcid.org/0000-0003-2630-4339>
 Dorota Konopka-Postupolska  <https://orcid.org/0000-0002-3181-5478>
 Jaiana Malabarba  <https://orcid.org/0000-0002-1263-9026>
 Anja K. Meents  <https://orcid.org/0000-0003-2267-6952>
 Axel Mithöfer  <https://orcid.org/0000-0001-5229-6913>
 Ralf Oelmüller  <https://orcid.org/0000-0002-3878-0044>
 Edgar Peiter  <https://orcid.org/0000-0002-9104-3238>
 Julia Rachowka  <https://orcid.org/0000-0002-4148-9928>
 Michael Reichelt  <https://orcid.org/0000-0002-6691-6500>
 Sandra S. Scholz  <https://orcid.org/0000-0002-5389-5296>
 Katie A. Wilkins  <https://orcid.org/0000-0001-6513-856X>

References

- Alexandersson E, Saalbach G, Larsson C, Kjellbom P. 2004. Arabidopsis plasma membrane proteomics identifies components of transport, signal transduction and membrane trafficking. *Plant and Cell Physiology* 45: 1543–1556.
- Alonso C, Ramos-Cruz D, Becker C. 2019. The role of plant epigenetics in biotic interactions. *New Phytologist* 221: 731–737.
- Arimura G, Köpke S, Kunert M, Volpe V, David A, Brand P, Dabrowska P, Maffei ME, Boland W. 2008. Effects of feeding *Spodoptera littoralis* on lima bean leaves: IV. Diurnal and nocturnal damage differentially initiate plant volatile emission. *Plant Physiology* 146: 965–973.
- Arimura GI, Ozawa R, Maffei ME. 2011. Recent advances in plant early signaling in response to herbivory. *International Journal of Molecular Sciences* 12: 3723–3739.
- Bergomaz R, Boppré M. 1986. A simple instant diet for rearing Arctiidae and other moths. *Journal of the Lepidopterists' Society* 40: 131–137.
- Bricchi I, Leitner M, Foti M, Mithöfer A, Boland W, Maffei ME. 2010. Robotic mechanical wounding (MecWorm) versus herbivore-induced responses: early signaling and volatile emission in lima bean (*Phaseolus lunatus* L.). *Planta* 232: 719–729.
- Carella P, Merl-Pham J, Wilson DC, Dey S, Hauck SM, Vlot C, Cameron RK. 2016. Comparative proteomics analysis of Arabidopsis phloem exudates collected during the induction of systemic acquired resistance. *Plant Physiology* 171: 149–1510.
- Clark G, Konopka-Postupolska D, Hennig J, Roux S. 2010. Is annexin 1 a multifunctional protein during stress responses? *Plant Signaling and Behavior* 5: 303–307.

- Clark GB, Morgan RO, Fernandez MP, Roux SJ. 2012. Evolutionary adaptation of plant annexins has diversified their molecular structures, interactions and functional roles. *New Phytologist* 196: 695–712.
- Clark GB, Sessions A, Eastburn DJ, Roux SJ. 2001. Differential expression of members of the annexin multigene family in Arabidopsis. *Plant Physiology* 126: 1072–1084.
- Davies J. 2014. Annexin-mediated calcium signalling in plants. *Plants* 3: 128–140.
- DeFalco TA, Bender KW, Snedden WA. 2010. Breaking the code: Ca²⁺ sensors in plant signalling. *Biochemical Journal* 425: 27–40.
- Demidchik V, Maathuis FJM. 2007. Physiological roles of nonselective cation channels in plants: from salt stress to signalling and development. *New Phytologist* 175: 387–404.
- Dengler NG. 2006. The shoot apical meristem and development of vascular architecture. *Canadian Journal of Botany* 84: 1660–1671.
- Dinneny JR, Long TA, Wang JY, Jung JW, Mace D, Pointer S, Barron C, Brady SM, Schiefelbein J, Benfey PN. 2008. Cell identity mediates the response of Arabidopsis roots to abiotic stress. *Science* 320: 942–945.
- Dodd AN, Kudla J, Sanders D. 2010. The language of calcium signaling. *Annual Review of Plant Biology* 61: 593–620.
- Espinoza C, Liang Y, Stacey G. 2017. Chitin receptor CERK1 links salt stress and chitin-triggered innate immunity in Arabidopsis. *Plant Journal* 89: 984–995.
- Farmer E, Mousavi S, Lenglet A. 2013. Leaf numbering for experiments on long distance signalling in Arabidopsis. *Protocol Exchange* doi: 10.1038/protex.2013.071.
- Fisahn J, Herde O, Willmitzer L, Peña-Cortés H. 2004. Analysis of the transient increase in cytosolic Ca²⁺ during the action potential of higher plants with high temporal resolution: requirement of Ca²⁺ transients for induction of jasmonic acid biosynthesis and PINII gene expression. *Plant and Cell Physiology* 45: 456–459.
- Foreman J, Demidchik V, Bothwell JHF, Mylona P, Miedema H, Torres MA, Linstead P, Costa S, Brownlee C, Jones JDG *et al.* 2003. Reactive oxygen species produced by NADPH oxidase regulate plant cell growth. *Nature* 422: 442–446.
- Guelette BS, Benning UF, Hoffmann-Benning S. 2012. Identification of lipids and lipid-binding proteins in phloem exudates from Arabidopsis thaliana. *Journal of Experimental Botany* 63: 3603–3616.
- Gully K, Pelletier S, Guillou M, Ferrand M, Aligon S, Pokotylo I, Perrin A, Vergne E, Fagard M, Ruelland E *et al.* 2019. The SCOOP12 peptide regulates defense response and root elongation in Arabidopsis thaliana. *Journal of Experimental Botany* 70: 1349–1365.
- Heyer M, Reichelt M, Mithöfer A. 2018a. A holistic approach to analyze systemic jasmonate accumulation in individual leaves of Arabidopsis rosettes upon wounding. *Frontiers in Plant Science* 871: e1569.
- Heyer M, Scholz SS, Voigt D, Reichelt M, Aldon D, Oelmüller R, Boland W, Mithöfer A. 2018b. Herbivory-responsive calmodulin-like protein CML9 does not guide jasmonate-mediated defenses in Arabidopsis thaliana. *PLoS ONE* 13: e0197633.
- Jami SK, Dalal A, Divya K, Kirti PB. 2009. Molecular cloning and characterization of five annexin genes from Indian mustard (*Brassica juncea* L. Czern and Coss). *Plant Physiology and Biochemistry* 47: 977–990.
- Kiep V, Vadassery J, Lattke J, Maaß JP, Boland W, Peiter E, Mithöfer A. 2015. Systemic cytosolic Ca²⁺ elevation is activated upon wounding and herbivory in Arabidopsis. *New Phytologist* 207: 996–1004.
- Knight H, Trewavas AJ, Knight MR. 1996. Cold calcium signaling in Arabidopsis involves two cellular pools and a change in calcium signature after acclimation. *Plant Cell* 8: 489–503.
- Knight MR, Campbell AK, Smith SM, Trewavas AJ. 1991. Transgenic plant aequorin reports the effects of touch and cold-shock and elicitors on cytoplasmic calcium. *Nature* 352: 524–526.
- Konopka-Postupolska D, Clark G, Goch G, Debski J, Floras K, Cantero A, Fijolek B, Roux S, Hennig J. 2009. The role of annexin 1 in drought stress in Arabidopsis. *Plant Physiology* 150: 1394–1410.
- Kovács I, Ayaydin F, Oberschall A, Ipacs L, Bottka S, Pongor S, Dudits D, Tóth ÉC. 1998. Immunolocalization of a novel annexin-like protein encoded by a stress and abscisic acid responsive gene in alfalfa. *The Plant Journal* 15: 185–197.
- Kudla J, Batistic O, Hashimoto K. 2010. Calcium signals: the lead currency of plant information processing. *Plant Cell* 22: 541–563.
- Kumari A, Chételat A, Nguyen CT, Farmer EE. 2019. Arabidopsis H⁺-ATPase AHA1 controls slow wave potential duration and wound-response jasmonate pathway activation. *Proceedings of the National Academy of Sciences, USA* 116: 20226–20231.
- Laohavisit A, Davies JM. 2009. Multifunctional annexins. *Plant Science* 177: 532–539.
- Laohavisit A, Davies JM. 2011. Annexins. *New Phytologist* 189: 40–53.
- Laohavisit A, Mortimer JC, Demidchik V, Coxon KM, Stancombe MA, Macpherson N, Brownlee C, Hofmann A, Webb AAR, Miedema H *et al.* 2009. *Zea mays* annexins modulate cytosolic free Ca²⁺ and generate a Ca²⁺-permeable conductance. *Plant Cell* 21: 479–493.
- Laohavisit A, Shang Z, Rubio L, Cuin TA, Véry A-A, Wang A, Mortimer JC, Macpherson N, Coxon KM, Battey NH *et al.* 2012. Arabidopsis annexin1 mediates the radical-activated plasma membrane Ca²⁺ and K⁺-permeable conductance in root cells. *Plant Cell* 24: 1522–1533.
- Lee S, Lee EJ, Yang EJ, Lee JE, Park AR, Song WH, Park OK. 2004. Proteomic identification of annexins, calcium-dependent membrane binding proteins that mediate osmotic stress and abscisic acid signal transduction in Arabidopsis. *Plant Cell* 16: 1378–1391.
- Lu Y, Ouyang B, Zhang J, Wang T, Lu C, Han Q, Zhao S, Ye Z, Li H. 2012. Genomic organization, phylogenetic comparison and expression profiles of annexin gene family in tomato (*Solanum lycopersicum*). *Gene* 499: 14–24.
- Ma L, Ye J, Yang Y, Lin H, Yue L, Luo J, Long Y, Fu H, Liu X, Zhang Y *et al.* 2019. The SOS2-SCaBP8 complex generates and fine-tunes an AtANN4-dependent calcium signature under salt stress. *Developmental Cell* 48: 697–709.
- Maffei ME, Arimura G-I, Mithöfer A. 2012. Natural elicitors, effectors and modulators of plant responses. *Natural Product Reports* 29: e1288.
- Maffei M, Bossi S, Spitter D, Mithöfer A, Boland W. 2004. Effects of feeding *Spodoptera littoralis* on lima bean leaves. I. Membrane potentials, intracellular calcium variations, oral secretions, and regurgitate components. *Plant Physiology* 134: 1752–1762.
- Maffei ME, Mithöfer A, Boland W. 2007. Before gene expression: early events in plant–insect interaction. *Trends in Plant Science* 12: 310–316.
- Maischak H, Zimmermann MR, Felle HH, Boland W, Mithöfer A. 2010. Alamehcin-induced electrical long distance signaling in plants. *Plant Signaling and Behavior* 5: 988–990.
- Marmagne A, Ferro M, Meinel T, Bruley C, Kuhn L, Garin J, Barbier-Brygoo H, Ephritikhine G. 2007. A high content in lipid-modified peripheral proteins and integral receptor kinases features in the Arabidopsis plasma membrane proteome. *Molecular & Cellular Proteomics* 6: 1980–1996.
- Meena MK, Prajapati R, Krishna D, Divakaran K, Pandey Y, Reichelt M, Mathew MKK, Boland W, Mithöfer A, Vadassery J. 2019. The Ca²⁺ channel CNGC19 regulates Arabidopsis defense against *Spodoptera* herbivory. *Plant Cell* 31: 1539–1562.
- Mithöfer A, Boland W. 2008. Recognition of herbivory-associated molecular patterns. *Plant Physiology* 146: 825–831.
- Mithöfer A, Boland W. 2012. Plant defense against herbivores: chemical aspects. *Annual Review of Plant Biology* 63: 431–450.
- Mithöfer A, Wanner G, Boland W. 2005. Effects of feeding *Spodoptera littoralis* on lima bean leaves. II. Continuous mechanical wounding resembling insect feeding is sufficient to elicit herbivory-related volatile emission. *Plant Physiology* 137: 1160–1168.
- Mohanta TK, Yadav D, Khan AL, Hashem A, Abd_Allah E, Al-Harrasi A. 2019. Molecular players of EF-hand containing calcium signaling event in plants. *International Journal of Molecular Sciences* 20: e1476.
- Mousavi SAR, Chauvin A, Pascaud F, Kellenberger S, Farmer EE. 2013. GLUTAMATE RECEPTOR-LIKE genes mediate leaf-to-leaf wound signalling. *Nature* 500: 422–426.
- Nguyen CT, Kurenda A, Stolz S, Chételat A, Farmer EE. 2018. Identification of cell populations necessary for leaf-to-leaf electrical signaling in a wounded plant. *Proceedings of the National Academy of Sciences, USA* 115: 10178–10183.

- Patel N, Hamamouch N, Li C, Hewezi T, Hussey RS, Baum TJ, Mitchum MG, Davis EL. 2010. A nematode effector protein similar to annexins in host plants. *Journal of Experimental Botany* 61: 235–248.
- Peiter E. 2011. The plant vacuole: emitter and receiver of calcium signals. *Cell Calcium* 50: 120–128.
- Peiter E. 2016. The ever-closer union of signals: propagating waves of calcium and ROS are inextricably linked. *Plant Physiology* 172: 3–4.
- Peiter E, Maathuis FJM, Mills LN, Knight H, Pelloux J, Hetherington AM, Sanders D. 2005. The vacuolar Ca^{2+} -activated channel TPC1 regulates germination and stomatal movement. *Nature* 434: 404–408.
- Pfaffl MW. 2001. A new mathematical model for relative quantification in real-time RT-PCR. *Nucleic Acids Research* 29: e45.
- Ranjan A, Vadassery J, Patel HK, Pandey A, Palaparthi R, Mithöfer A, Sonti RV. 2015. Upregulation of jasmonate biosynthesis and jasmonate-responsive genes in rice leaves in response to a bacterial pathogen mimic. *Functional and Integrative Genomics* 15: 363–373.
- Richards SL, Laohavisit A, Mortimer JC, Shabala L, Swarbreck SM, Shabala S, Davies JM. 2014. Annexin 1 regulates the H_2O_2 -induced calcium signature in *Arabidopsis thaliana* roots. *The Plant Journal* 77: 136–145.
- Ruijter JM, Ramakers C, Hoogaars WMH, Karlen Y, Bakker O, van den Hoff MJB, Moorman AFM. 2009. Amplification efficiency: linking baseline and bias in the analysis of quantitative PCR data. *Nucleic Acids Research* 37: e45.
- Saijo Y, Loo EP. 2019. Plant immunity in signal integration between biotic and abiotic stress responses. *New Phytologist* 225: 87–104.
- Scholz SS, Heyer M, Vadassery J, Mithöfer A. 2016. A role for calmodulin-like proteins in herbivore defense pathways in plants. *Journal of Endocytobiosis and Cell Research* 27: 1–12.
- Scholz SS, Malabarba J, Reichelt M, Heyer M, Ludewig F, Mithöfer A. 2017. Evidence for GABA-induced systemic GABA accumulation in *Arabidopsis* upon wounding. *Frontiers in Plant Science* 8: e388.
- Scholz SS, Vadassery J, Heyer M, Reichelt M, Bender KW, Snedden WA, Boland W, Mithöfer A. 2014. Mutation of the *Arabidopsis* calmodulin-like protein CML37 deregulates the jasmonate pathway and enhances susceptibility to herbivory. *Molecular Plant* 7: 1712–1726.
- Seybold H, Trempe F, Ranf S, Scheel D, Romeis T, Lee J. 2014. Ca^{2+} signalling in plant immune response: from pattern recognition receptors to Ca^{2+} decoding mechanisms. *New Phytologist* 204: 782–790.
- Sutter J, Sieben C, Hartel A, Eisenach C, Thiel G, Blatt MR. 2007. Abscisic acid triggers the endocytosis of the *Arabidopsis* KAT1 K^+ channel and its recycling to the plasma membrane. *Current Biology* 17: 1396–1402.
- Swarbreck SM, Colaco R, Davies JM. 2013. Plant calcium-permeable channels. *Plant Physiology* 163: 514–522.
- Tai L, Li B-B, Nie X, Zhang P-P, Hu C, Zhang L, Liu W-T, Li W-Q, Chen K. 2019. Calmodulin is the fundamental regulator of NADK-mediated NAD signaling in plants. *Frontiers in Plant Science* 10: e681.
- Toyota M, Spencer D, Sawai-Toyota S, Jiaqi W, Zhang T, Koo AJ, Howe GA, Gilroy S. 2018. Glutamate triggers long-distance, calcium-based plant defense signaling. *Science* 361: 1112–1115.
- Vadassery J, Reichelt M, Hause B, Gershenzon J, Boland W, Mithöfer A. 2012a. CML42-mediated calcium signaling coordinates responses to *Spodoptera* herbivory and abiotic stresses in *Arabidopsis*. *Plant Physiology* 159: 1159–1175.
- Vadassery J, Reichelt M, Jimenez-Aleman GH, Boland W, Mithöfer A. 2014. Neomycin inhibition of (+)-7-*iso*-jasmonoyl-L-isoleucine accumulation and signaling. *Journal of Chemical Ecology* 40: 676–686.
- Vadassery J, Scholz SS, Mithöfer A. 2012b. Multiple calmodulin-like proteins in *Arabidopsis* are induced by insect-derived (*Spodoptera littoralis*) oral secretion. *Plant Signaling and Behavior* 7: 1277–1280.
- Verrillo F, Occhipinti A, Kanchiswamy CN, Maffei ME. 2014. Quantitative analysis of herbivore-induced cytosolic calcium by using a Cameleon (YC 3.6) calcium sensor in *Arabidopsis thaliana*. *Journal of Plant Physiology* 171: 136–139.
- Vincent TR, Avramova M, Canham J, Higgins P, Bilkey N, Mugford ST, Pitino M, Toyota M, Gilroy S, Miller AJ *et al.* 2017. Interplay of plasma membrane and vacuolar ion channels, together with BAK1, elicits rapid cytosolic calcium elevations in *Arabidopsis* during aphid feeding. *Plant Cell* 29: 1460–1479.
- Wasternack C. 2007. Jasmonates: an update on biosynthesis, signal transduction and action in plant stress response, growth and development. *Annals of Botany* 100: 681–697.
- Xu L, Tang Y, Gao S, Su S, Hong L, Wang W, Fang Z, Li X, Ma J, Quan W *et al.* 2016. Comprehensive analyses of the annexin gene family in wheat. *BMC Genomics* 17: e415.
- Yan C, Fan M, Yang M, Zhao J, Zhang W, Su Y, Xiao L, Deng H, Xie D. 2018. Injury activates Ca^{2+} /calmodulin-dependent phosphorylation of JAV1-JAZ8-WRKY51 complex for jasmonate biosynthesis. *Molecular Cell* 70: 136–149.
- Yuan J, Wen Z, Gu C, Wang D. 2014. Introduction of high throughput and cost effective SNP genotyping platforms in soybean. *Plant Genetics, Genomics, and Biotechnology* 2: 90–94.
- Zebelo S, Piorkowski J, Disi J, Fadamiro H. 2014. Secretions from the ventral eversible gland of *Spodoptera exigua* caterpillars activate defense-related genes and induce emission of volatile organic compounds in tomato, *Solanum lycopersicum*. *BMC Plant Biology* 14: e140.
- Zhao J, Li L, Liu Q, Liu P, Li S, Yang D, Chen Y, Pagnotta S, Favery B, Abad P *et al.* 2019. A MIF-like effector suppresses plant immunity and facilitates nematode parasitism by interacting with plant annexins. *Journal of Experimental Botany* 70: 5943–5958.
- Zimmermann MR, Maischak H, Mithöfer A, Boland W, Felle HH. 2009. System potentials, a novel electrical long-distance apoplastic signal in plants, induced by wounding. *Plant Physiology* 149: 1593–1600.

Supporting Information

Additional Supporting Information may be found online in the Supporting Information section at the end of the article.

Fig. S1 $[Ca^{2+}]_{cyt}$ response upon mechanical wounding (MW) and water (H_2O) in Col-0 and annexin1 mutants.

Fig. S2 Weight of *Spodoptera littoralis* larvae fed on *Arabidopsis* Col-0 and *ann1-1* plants expressing *Aequorin*.

Fig. S3 Accumulation of related jasmonates after wounding and oral secretion (OS) treatment.

Video S1 Time-lapse of $[Ca^{2+}]_{cyt}$ response in *Arabidopsis thaliana* Col-0 rosette induced by mechanical wounding + oral secretion (OS) of the leaf lamina.

Video S2 Time-lapse of $[Ca^{2+}]_{cyt}$ response in *Arabidopsis thaliana ann1-1* rosette induced by mechanical wounding + oral secretion (OS) of the leaf lamina.

Video S3 Time-lapse of $[Ca^{2+}]_{cyt}$ response in *Arabidopsis thaliana* Col-0 rosette induced by mechanical wounding + water of the leaf lamina.

Video S4 Time-lapse of $[Ca^{2+}]_{cyt}$ response in *Arabidopsis thaliana ann1-1* rosette induced by mechanical wounding + water of the leaf lamina.

Please note: Wiley Blackwell are not responsible for the content or functionality of any Supporting Information supplied by the authors. Any queries (other than missing material) should be directed to the *New Phytologist* Central Office.