

Supplementary material

Non-specific clustering of Histidine tagged Green Fluorescent Protein mediated by surface interactions: the collective effect in the protein-adsorption behaviour

Tomasz D. Sobieściak* and Piotr Zielenkiewicz


*Institute of Biochemistry and Biophysics, Polish Academy of Sciences, Pawinskiego 5a, 02-106
Warsaw, Poland*

* Author to whom correspondence should be addressed. e-mail: tomasz_s@ibb.waw.pl

Fluorescence Microphotographs

Figure S1. Micropatterned surfaces functionalized with NTA after exposure to the His-tagged GFP. The same fragment of chip observed under increased microscope magnifications.

a) NTA-functionalized internal area of squares loaded with nickel(II) ions. Samples were incubated overnight at ambient temperature and subsequently washed with PBS buffer.



b) Micropatterned surface after washing with buffer containing 250 mM imidazole (after 1 day of incubation and subsequent washing step).






Figure S2. Micropatterned surfaces functionalized with tacn-bis(formyl) after exposure to the His-tagged GFP. The same fragment of chip observed under increased microscope magnifications.


a) Tacn-bis(formyl)-functionalized internal area of squares after loading with nickel ions. Samples were incubated overnight at ambient temperature and subsequently washed with PBS buffer.



b) Micropatterned surface after washing with buffer containing 250 mM imidazole (after 1 day of incubation and subsequent washing step).



c) Micropatterned surface after washing with buffer containing 200 mM EDTA (after 5 min of incubation and subsequent washing step).



d) His₆-GFP deposited on the micropatterned surface presenting tacn-bis(formyl) without nickel ions (after overnight incubation and subsequent washing step).




Figure S3. Sequence of microphotographs – from 0 s to 60 s after loading of His-GFP on a whole surface of chip functionalized with tacn-bis(formyl) (without nickel ions) in internal area of squares. Protein is deposited directly under fluorescent microscope objective and observed immediately. Images are arranged in a series of pictures taken at a specific time (magnification 100x).






Figure S4. Micropatterned surfaces functionalized with N-hydroxysuccinimide ester after exposure to GFP or to fluorescently labeled proteins. The same fragment of chip observed under increased microscope magnifications.

a) His-GFP covalently bound to the micropatterned surface; images were acquired after overnight incubation and subsequent washing of chips with PBS buffer.



b) FITC–Con A covalently bound to the micropatterned surface; images were acquired after overnight incubation and subsequent washing of chips with PBS buffer.



c) Texas Red labeled BSA covalently bound to the micropatterned surface; images were acquired after overnight incubation and subsequent washing of chips with PBS buffer.

